

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 December 2003 (04.12.2003)

PCT

(10) International Publication Number
WO 03/099818 A1

(51) International Patent Classification⁷: **C07D 471/04**,
475/04, 487/04, A61K 31/517, 31/519, A61P 3/10

(74) Agents: **FRIEDRICHSEN, Bernard, P. et al.**; Foley &
Lardner, 150 East Gilman Street, P.O. Box 1497, Madison,
WI 53701-1497 (US).

(21) International Application Number: **PCT/US03/16442**

(22) International Filing Date: **23 May 2003 (23.05.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/382,762 23 May 2002 (23.05.2002) US
60/441,019 17 January 2003 (17.01.2003) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **CHI-
RON CORPORATION** [US/US]; 4560 Horton Street,
Emeryville, CA 94608-2917 (US).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **BOYCE, Rustum**,
S. [IN/US]; 1818 Broadway, Apartment 206, San Fran-
cisco, CA 94109 (US). **AURRECOECHEA, Natalia**
[ES/US]; 3040 Brookfield Avenue, Oakland, CA 94605
(US). **CHU, Daniel** [US/US]; 3767 Benton Street, Santa
Clara, CA 95051 (US). **SMITH, Aaron** [US/US]; 2974
Flint Street, Union City, CA 94587 (US).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*


(54) Title: **SUBSTITUTED QUINAZOLINONE COMPOUNDS**



(IA)



(IB)



(IC)

(57) Abstract: A variety of low molecular weight, guanidino-containing molecules capable of acting as MC4-R agonists are pro-
vided. The compounds are useful in treating MC4-R mediated diseases. The compounds have the structure of Formulas (IA), (IB),
or (IC): where the values of the variable are defined herein.

WO 03/099818 A1

SUBSTITUTED QUINAZOLINONE COMPOUNDS

Field of the Invention

[0001] This invention relates to melanocortin-4 receptor (MC4-R) agonists and methods of their preparation. The invention also relates to methods of treating melanocortin-4 receptor-mediated diseases, such as obesity or diabetes, by activating the melanocortin-4 receptor with compounds provided herein.

Background of the Invention

[0002] Melanocortins are peptide products resulting from post-translational processing of pro-opiomelanocortin and are known to have a broad array of physiological activities. The natural melanocortins include the different types of melanocyte stimulating hormone (α -MSH, β -MSH, γ -MSH) and ACTH. Of these, α -MSH and ACTH are considered to be the main endogenous melanocortins.

[0003] The melanocortins mediate their effects through melanocortin receptors (MC-Rs), a subfamily of G-protein coupled receptors. There are at least five different receptor subtypes (MC1-R to MC5-R). MC1-R mediates pigmentation of the hair and skin. MC2-R mediates the effects of ACTH on steroidogenesis in the adrenal gland. MC3-R and MC4-R are predominantly expressed in the brain. MC5-R is considered to have a role in the exocrine gland system.

[0004] The melanocortin-4 receptor (MC4-R) is a seven-transmembrane receptor. MC4-R may participate in modulating the flow of visual and sensory information, coordinate aspects of somatomotor control, and/or participate in the modulation of autonomic outflow to the heart. K. G. Mountjoy *et al.*, *Science*, 257:1248-125 (1992). Significantly, inactivation of this receptor by gene targeting has resulted in mice that develop a maturity

onset obesity syndrome associated with hyperphagia, hyperinsulinemia, and hyperglycemia. D. Husznar *et al.*, *Cell*, 88(1): 131-41 (1997). MC4-R has also been implicated in other disease states including erectile disorders, cardiovascular disorders, neuronal injuries or disorders, inflammation, fever, cognitive disorders, and sexual behavior disorders. M. E. Hadley and C. Haskell-Luevano, The proopiomelanocortin system, *Ann. N. Y. Acad. Sci.*, 885:1 (1999).

[0005] Furthermore, observations in connection with endogenous MC4-R antagonists indicate that MC4-R is implicated in endogenous energy regulation. For example, an agouti protein is normally expressed in the skin and is an antagonist of the cutaneous MC receptor involved in pigmentation, MC1-R. M. M. Ollmann *et al.*, *Science*, 278:135-138 (1997). However, overexpression of agouti protein in mice leads to a yellow coat color due to antagonism of MC1-R and increased food intake and body weight due to antagonism of MC4-R. L. L. Kiefer *et al.*, *Biochemistry*, 36: 2084-2090 (1997); D. S. Lu *et al.*, *Nature*, 371:799-802 (1994). Agouti related protein (AGRP), an agouti protein homologue, antagonizes MC4-R but not MC1-R. T. M. Fong *et al.*, *Biochem. Biophys. Res. Commun.* 237:629-631 (1997). Administration of AGRP in mice increases food intake and causes obesity but does not alter pigmentation. M. Rossi *et al.*, *Endocrinology*, 139:4428-4431 (1998). Together, this research indicates that MC4-R participates in energy regulation, and therefore, identifies this receptor as a target for a rational drug design for the treatment of obesity.

[0006] In connection with MC4-R and its uncovered role in the etiology of obesity and food intake, the prior art includes reports of compounds and compositions that act as agonists or antagonists of MC4-R. As examples, U.S. Patent No. 6,060,589 describes polypeptides that are capable of modulating signaling activity of melanocortin receptors. Also, U.S. Patent Nos. 6,054,556 and 5,731,408 describe families of agonists and antagonists for MC4-R receptors that are lactam heptapeptides having a cyclic structure. WO 01/10842 discloses MC4-R binding compounds having a multitude of

structures and methods of using such compounds to treat MC4-R associated disorders. Some of the compounds described include amidino- and guanidino-containing arenes and heteroarenes.

[0007] Various other classes of compounds have been disclosed as having MC4-R agonist activity. For example, WO 01/70708 and WO 00/74679 disclose MC4-R agonists that are piperidine compounds and derivatives, while WO 01/70337 and WO 99/64002 disclose MC-R agonists that are spiro piperidine derivatives. Other known melanocortin receptor agonists include aromatic amine compounds containing amino acid residues, particularly tryptophan residues, as disclosed in WO 01/55106. Similar agonists are disclosed in WO 01/055107 which comprise aromatic amine compounds containing tertiary amide or tertiary amine groups. Finally, WO 01/055109 discloses melanocortin receptor agonists comprising aromatic amines which are generally bisamides separated by a nitrogen-containing alkyl linker.


[0008] Guanidine-containing compounds having a variety of biological activities are also known in the prior art. For example, U.S. patent No. 4,732,916 issued to Satoh *et al.* discloses guanidine compounds useful as antiulcer agents; U.S. Patent No. 4,874,864, U.S. Patent No. 4,949,891, and U.S. Patent No. 4,948,901 issued to Schnur *et al.* and EP 0343 894 disclose guanidino compounds useful as protease inhibitors and as anti-plasmin and anti-thrombin agents; and U.S. Patent No. 5,352,704 issued to Okuyama *et al.* discloses a guanidino compound useful as an antiviral agent. Guanidine-containing compounds are also disclosed in other references. For example, U.S. Patent No. 6,030,985 issued to Gentile *et al.* discloses guanidine compounds useful for treating and preventing conditions in which inhibition of nitric oxide synthetase is beneficial such as stroke, schizophrenia, anxiety, and pain. U.S. Patent No. 5,952,381 issued to Chen *et al.* discloses certain guanidine compounds for use in selectively inhibiting or antagonizing $\alpha_v\beta_3$ integrins.

[0009] Various 5-, 6-, and 7- membered fully saturated 1-azacarbocyclic-2-ylidene derivatives of guanidine are disclosed as having anti-secretory and hypoglycemic activities by U.S. Patent No. 4,211,867 issued to Rasmussen. Such compounds are also taught as useful for the treatment of cardiovascular disease. Other guanidine derivatives are disclosed by U.S. Patent No. 5,885,985 issued to Macdonald *et al.* as useful in therapy to treat inflammation. Various guanidinobenzamide compounds are disclosed in WO 02/18327. The guanidinobenzamides are disclosed as useful for treating obesity and type II diabetes.


[0010] Nevertheless, there remains a need for potent and specific agonists of MC4-R that are low molecular weight small molecules. Methods of treating a melanocortin-4 receptor mediated disease, such as obesity, with such non-peptide drugs, are also particularly desirable.

Summary of the Invention


[0011] The instant invention provides potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, there has been provided, in accordance with one aspect of the invention, compounds of formula IA, IB, or IC:



IA



IB



IC

wherein

Z^1 is selected from the group consisting of CR^4 and N;

Z^2 is selected from the group consisting of CR^5 and N;

Z^3 is selected from the group consisting of CR^6 and N;


R^1 is selected from the group consisting of substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;

R^2 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, alkylcarbonyl, and arylcarbonyl groups;


R^3 is selected from the group consisting of H, and substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;

R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclylamino, heteroaryl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl groups;

W is a group of formula IIA or IIB;



IIA



IIB

$R^{1'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

$R^{2'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

or $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

$R^{3'}$ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups; and

R^4 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocycl, arylalkyl, and heteroarylalkyl groups.

[0012] Compounds provided by the invention further include prodrugs of the compound of formula IA, IB, and IC, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0013] The invention further provides compounds of formula IA and IC in which R^3 is H.

[0014] The invention further provides compounds of formula IA, IB, and IC in which Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group. In some such embodiments, at least one of R^4 , R^5 , or R^6 is a F.

[0015] The invention further provides compounds of formula IA, IB, and IC in which at least one of Z^1 , Z^2 , or Z^3 is N. In some such embodiments, Z^1 is N. In other such embodiments, Z^2 is N.

[0016] The invention further provides compounds of formula IA, IB, and IC in which R^3 is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In other embodiments of compounds of formula IA, IB, and IC, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-

dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl, bornyl, norbornyl, and decalinyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups.

[0017] The invention further provides compounds of formula IA, IB, and IC in which R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl,

arylalkyl, and heteroarylalkyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R¹ and R² may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

[0018] In still other embodiments of compounds of formula IA, IB, and IC, R¹ and R², together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In still other embodiments of compounds of formula IA, IB, and IC, R¹ and R², together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R¹ and R² are bound. In still other embodiments of compounds of formula IA, IB, and IC, R¹ and R², together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still other embodiments of compounds of formula IA, IB, and IC, R¹ and R², together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two methyl groups.

[0019] The invention further provides compounds of formula IA, IB, and IC in which R¹ is a 2,4-disubstituted phenylethyl group. In still other embodiments of compounds of formula IA, IB, and IC, R¹ is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-dialkylphenylethyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R¹ is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-

methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroarylloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroarylloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups.


[0020] In still further embodiments, the invention provides compounds of formula IA, IB and IC in which R^1 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0021] The invention further provides compounds of formula IB in which R^2 is H.

[0022] In still further embodiments, the invention provides compounds of formula IA and IC in which R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in which R^3 is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R^3 is selected from substituted and unsubstituted 2-

pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0023] The instant invention provides a second group of potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, there has been provided, in accordance with one aspect of the invention, compounds of formula IA and IC:



wherein

Z^1 is selected from the group consisting of CR^4 and N;

Z^2 is selected from the group consisting of CR^5 and N;

Z^3 is selected from the group consisting of CR^6 and N;


R^1 is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;

R^3 is selected from the group consisting of substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, aryloxyalkyl, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkyl groups, and $-C(=NH)$ -heterocyclyl groups, and groups of formula $-LR^7$;


R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino,

cycloalkyl, heterocyclamino, heteroarylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclaminocarbonyl, and heteroarylamino carbonyl groups;

W is a group of formula IIA or IIB;



IIA



IIB

R^{1'} is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

R² is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;


or R^{1'} and R², together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

R³ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

R^{4'} is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl groups;

L is selected from the group consisting of a covalent bond, -CH₂-, -O-, -S-, and -NH-;

R⁷ is selected from the group consisting of substituted and unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from a group of formula IIC;



IIC

wherein Y is selected from the group consisting of CH₂, O, S, and NR⁹ where R⁹ is selected from the group consisting of H, and substituted or unsubstituted alkyl groups;

R⁸ is selected from the group consisting of H, a halogen, hydroxyl, carboxylic acid, and substituted or unsubstituted alkyl, amino, alkylamino, dialkylamino, alkylaminoalkyl, heterocyclyl, alkoxy, carbonyl, and aminocarbonyl groups; and

m is an integer selected from 0, 1, and 2.

[0024] In some aspects of the invention, compounds provided of the second group of potent and specific agonists of MC4-R further include prodrugs of the compounds of formula IA and IC, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0025] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R¹ is H or an alkyl group such as a methyl, ethyl, propyl, butyl, or pentyl group. In some such embodiments, R¹ is H whereas in other such embodiments, R¹ is a methyl group.

[0026] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group. In some such embodiments, at least one of R^4 , R^5 , or R^6 is a F.

[0027] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which at least one of Z^1 , Z^2 , or Z^3 is N. In some such embodiments, Z^1 is N. In other such embodiments, Z^2 is N.

[0028] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R^3 is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In other embodiments of the second group of compounds of formula IA and IC, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-

phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkylnorbornyl, bornyl, norbornyl, and decalynyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups.

[0029] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-

dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

[0030] In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R^1 and R^2 are bound. In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still other embodiments of the second group of compounds of formula IA and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two methyl groups.

[0031] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R^1 is a 2,4-disubstituted phenylethyl group. In still other embodiments of the second group of compounds of formula IA and IC, R^1 is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-dialkylphenylethyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^1 is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl,

2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups.

[0032] In still further embodiments, the invention provides compounds of formula IA and IC in which R^1 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0033] In still further embodiments, the invention provides compounds of formula IA and IC in which R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in which R^3 is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R^3 is selected from substituted and unsubstituted 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0034] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R³ is a 2,4-disubstituted phenylethyl group. In still other embodiments of the second group of compounds of formula IA and IC, R³ is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-dialkylphenylethyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R³ is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxyethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxyethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxyethyl)ethyl groups. In some of the embodiments described in this paragraph, R¹ is H or is an alkyl group having from one to eight carbon atoms. In some of the embodiments described in this paragraph,


R¹ is H whereas in other such embodiments R¹ is a methyl, ethyl, or propyl group.

[0035] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R³ is a substituted alkyl group such as a substituted or unsubstituted aryloxyalkyl group or a substituted or unsubstituted heteroaryloxyalkyl group. In some such embodiments, R³ is a substituted or unsubstituted aryloxymethyl group. In some such embodiments, of the second group of compounds of formula IA and IC, R³ is selected from the group consisting of -CH₂-O-aryl groups where the aryl group is substituted with one or more halogen group such as with one or more Cl or F. In some such embodiments, the aryl group is additionally substituted with an alkoxy group such as a methoxy or ethoxy group. In some embodiments of the second group of compounds of formula IA and IC, R³ is a -CH₂-O-aryl group where the aryl group is selected from the group consisting of 2,4-difluorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-methoxyphenyl, 2,4-dichlorophenyl, 4-chlorophenyl, 2-chlorophenyl, and 2-chloro-4-methoxyphenyl groups. In other embodiments of the second group of compounds of formula IA and IC, R³ is a heterocyclalkyl group. In some other embodiments of compounds of formula IA and IC, R³ is a substituted or unsubstituted arylalkoxyalkyl group or a heteroarylalkoxyalkyl group.

[0036] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R³ is a substituted or unsubstituted heterocyclalkyl group. In some such embodiments, R³ is a substituted or unsubstituted heterocyclmethyl group. In some such embodiments, the heterocycl group is selected from the group consisting of substituted and unsubstituted 1*H*-tetrazole, piperazine, piperidine, imidazole, and morpholine groups. In some such embodiments, R³ is a -CH₂-heterocycl group where the heterocycl group is a 1*H*-tetrazole, an imidazole, an N-methylpiperazine, a 4-hydroxypiperidine, a 3-hydroxypiperidine, or a morpholine. In still other

embodiments of the second group of compounds of formula IA and IC, R^3 is a heterocyclyl group. In some embodiments of the second group of compounds of formula IA and IC, R^3 is a substituted or unsubstituted piperazinyl group such as an N-methylpiperazinyl group, is a substituted or unsubstituted pyridine group, is a substituted or unsubstituted tetrazole group, is a substituted or unsubstituted cycloalkyl group such as a 4-methylcyclohexyl group, or is a substituted or unsubstituted phenyl group. In yet other embodiments of the second group of compounds of formula IA and IC, R^3 is an alkoxyalkyl group such as a methoxyalkyl group or an ethoxyalkyl group. In some such embodiments, R^3 is an alkoxyalkyl group such as an alkoxymethyl group such as a methoxymethyl group.

[0037] The instant invention provides a third group of potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, there has been provided, in accordance with one aspect of the invention, compounds of formula ID:



wherein


R^1 is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;

R^2 and R^3 are independently selected from the group consisting of -H, substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, aryloxyalkyl, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl,


cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkyl groups, and $-C(=NH)$ -heterocycl groups, and groups of formula $-LR^7$;

R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclamino, heteroaryl amino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclaminocarbonyl, and heteroarylaminocarbonyl groups;

W is a group of formula IIA or IIB;



IIA



IIB

$R^{1'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocycl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

$R^{2'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocycl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

or $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocycl or heteroaryl group;


$R^{3'}$ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,

heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

R^4 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl groups;

L is selected from the group consisting of a covalent bond, $-CH_2-$, $-O-$, $-S-$, and $-NH-$;

R^7 is selected from the group consisting of substituted and unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from a group of formula IIC;



IIC

wherein Y is selected from the group consisting of CH_2 , O, S, and NR^9 where R^9 is selected from the group consisting of H, and substituted or unsubstituted alkyl groups;

R^8 is selected from the group consisting of H, a halogen, hydroxyl, carboxylic acid, and substituted or unsubstituted alkyl, amino, alkylamino, dialkylamino, alkylaminoalkyl, heterocyclyl, alkoxy, carbonyl, and aminocarbonyl groups; and

m is an integer selected from 0, 1, and 2.

[0038] In some aspects of the invention, compounds provided of the third group of potent and specific agonists of MC4-R further include prodrugs of the compounds of formula ID, pharmaceutically acceptable salts thereof,

stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0039] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^1 is H or an alkyl group such as a methyl, ethyl, propyl, butyl, or pentyl group. In some such embodiments, R^1 is H whereas in other such embodiments, R^1 is a methyl group.

[0040] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^2 is H or an alkyl group such as a methyl, ethyl, propyl, butyl, or pentyl group.

[0041] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which at least one of R^4 , R^5 , or R^6 is a F.

[0042] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which $R^{3'}$ is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In other embodiments of the third group of compounds of formula ID, $R^{3'}$ is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-

dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl, bornyl, norbornyl, and decalanyl groups. In still other embodiments of the third group of compounds of formula ID, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups.

[0043] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the third group of compounds of formula ID, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 may be the same or different

and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

[0044] In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R^1 and R^2 are bound. In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still other embodiments of the third group of compounds of formula ID, R^1 and R^2 , together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two methyl groups.

[0045] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^1 is a 2,4-disubstituted phenylethyl group. In still other embodiments of the third group of compounds of formula ID, R^1 is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-dialkylphenylethyl groups. In still other embodiments of the third group of compounds of formula ID, R^1 is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-

methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups.

[0046] In still further embodiments, the invention provides compounds of formula ID in which R^1 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0047] In still further embodiments, the invention provides compounds of formula ID in which R^2 is H or a substituted or unsubstituted alkyl group.

[0048] In still further embodiments, the invention provides compounds of formula ID in which R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in

which R³ is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R³ is selected from substituted and unsubstituted 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0049] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R³ is a 2,4-disubstituted phenylethyl group. In still other embodiments of the third group of compounds of formula ID, R³ is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-dialkylphenylethyl groups. In still other embodiments of the third group of compounds of formula ID, R³ is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted


(heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups. In some of the embodiments described in this paragraph, R^1 is H or is an alkyl group having from one to eight carbon atoms. In some of the embodiments described in this paragraph, R^1 is H whereas in other such embodiments R^1 is a methyl, ethyl, or propyl group.

[0050] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^3 is a substituted alkyl group such as a substituted or unsubstituted aryloxyalkyl group or a substituted or unsubstituted heteroaryloxyalkyl group. In some such embodiments, R^3 is a substituted or unsubstituted aryloxymethyl group. In some such embodiments, of the third group of compounds of formula ID, R^3 is selected from the group consisting of $-CH_2-O$ -aryl groups where the aryl group is substituted with one or more halogen group such as with one or more Cl or F. In some such embodiments, the aryl group is additionally substituted with an alkoxy group such as a methoxy or ethoxy group. In some embodiments of the third group of compounds of formula ID, R^3 is a $-CH_2-O$ -aryl group where the aryl group is selected from the group consisting of 2,4-difluorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-methoxyphenyl, 2,4-dichlorophenyl, 4-chlorophenyl, 2-chlorophenyl, and 2-chloro-4-methoxyphenyl groups. In other embodiments of the third group of compounds of formula ID, R^3 is a heterocyclylalkyl group. In some other embodiments of compounds of formula ID, R^3 is a substituted or unsubstituted arylalkoxyalkyl group or a heteroarylalkoxyalkyl group.

[0051] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^3 is a substituted or unsubstituted heterocyclylalkyl group. In some such embodiments, R^3 is a substituted or

unsubstituted heterocyclymethyl group. In some such embodiments, the heterocyclyl group is selected from the group consisting of substituted and unsubstituted 1*H*-tetrazole, piperazine, piperidine, imidazole, and morpholine groups. In some such embodiments, R³ is a -CH₂-heterocyclyl group where the heterocyclyl group is a 1*H*-tetrazole, an imidazole, an N-methylpiperazine, a 4-hydroxypiperidine, a 3-hydroxypiperidine, or a morpholine. In still other embodiments of the third group of compounds of formula ID, R³ is a heterocyclyl group. In some embodiments of the third group of compounds of formula ID, R³ is a substituted or unsubstituted piperazinyl group such as an N-methylpiperazinyl group, is a substituted or unsubstituted pyridine group, is a substituted or unsubstituted tetrazole group, is a substituted or unsubstituted cycloalkyl group such as a 4-methylcyclohexyl group, or is a substituted or unsubstituted phenyl group. In yet other embodiments of the third group of compounds of formula ID, R³ is an alkoxyalkyl group such as a methoxyalkyl group or an ethoxyalkyl group. In some such embodiments, R³ is a an alkoxyalkyl group such as an alkoxymethyl group such as a methoxymethyl group.

[0052] The instant invention provides a fourth group of potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, there has been provided, in accordance with one aspect of the invention, compounds of formula IE:




wherein


R^1 is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;

R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclalkyl, heteroarylalkyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclalkylaminocarbonyl, and heteroarylaminocarbonyl groups;

W is a group of formula IIA or IIB;



IIA



IIB

R^1 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

R^2 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

or R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

R^3 is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,

heterocyclyl, heterocyclalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups; and

$R^{4'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl groups.

[0053] In some aspects of the invention, compounds provided of the fourth group of potent and specific agonists of MC4-R further include prodrugs of the compounds of formula IE, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0054] In some aspects of the invention of compounds of the fourth group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IE in which R^1 is H or an alkyl group such as a methyl, ethyl, propyl, butyl, or pentyl group. In some such embodiments, R^1 is H whereas in other such embodiments, R^1 is a methyl group.

[0055] In some aspects of the invention of compounds of the fourth group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IE in which at least one of R^4 , R^5 , or R^6 is a F.

[0056] In some aspects of the invention of compounds of the fourth group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IE in which R^3 is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In other embodiments of the fourth group of compounds of formula IE, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-

aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl, bornyl, norbornyl, and decalinyl groups. In still other embodiments of the fourth group of compounds of formula IE, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups.

[0057] In some aspects of the invention of compounds of the fourth group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IE in which R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the fourth group of compounds of

formula IE, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

[0058] In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R^1 and R^2 are bound. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two methyl groups.

[0059] In some aspects of the invention of compounds of the fourth group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IE in which R^1 is a 2,4-disubstituted phenylethyl group. In still other embodiments of the fourth group of compounds of formula IE, R^1 is selected from the group consisting of 2,4-dihalophenylethyl, and 2,4-

dialkylphenylethyl groups. In still other embodiments of the fourth group of compounds of formula IE, R¹ is selected from the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups.

[0060] In still further embodiments, the invention provides compounds of formula IE in which R¹ is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0061] There has also been provided, in accordance with another aspect of the invention, a composition such as a pharmaceutical formulation

or medicament comprising a compound according to the instant invention and a pharmaceutically acceptable carrier. The invention further provides the use of the compounds of the invention in preparing a medicament for use in treating an MC4-R mediated disease. In some embodiments, such a disease is obesity or type II diabetes.

[0062] There has also been provided, in accordance with another aspect of the invention, a method of treating an MC4-R mediated disease, comprising administering to a subject in need thereof, a compound or composition of the instant invention.

[0063] In one embodiment, a disease to be treated by those methods of the instant invention is obesity or type II diabetes.

[0064] In one embodiment, a compound or composition of the invention is intranasally administered.

[0065] In one embodiment, a compound or composition of the invention is administered to a human subject.

[0066] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

Detailed Description of the Preferred Embodiment

[0067] The instant invention relates to novel classes of small molecule melanocortin-4 receptor (MC4-R) agonists. These compounds can be formulated into compositions and are useful in activating MC4-R, or in the treatment of MC4-R-mediated diseases, such as obesity, type II diabetes,

erectile dysfunction, polycystic ovary disease, complications resulting from or associated with obesity and diabetes, and Syndrome X.

[0068] The following definitions are used throughout this specification.

[0069] Alkyl groups include straight chain and branched alkyl groups having from 1 to about 8 carbon atoms. Examples of straight chain alkyl groups include methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, and octyl groups. Examples of branched alkyl groups, include, but are not limited to, isopropyl, sec-butyl, t-butyl, and isopentyl groups. Representative substituted alkyl groups may be substituted one or more times with, for example, amino, thio, alkoxy, or halo groups such as F, Cl, Br, and I groups.

[0070] Cycloalkyl groups are cyclic alkyl groups such as, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl groups. Cycloalkyl groups also includes rings that are substituted with straight or branched chain alkyl groups as defined above, and further include cycloalkyl groups that are substituted with other rings including fused rings such as, but not limited to, decalinyl, tetrahydronaphthyl, and indanyl. Cycloalkyl groups also include polycyclic cycloalkyl groups such as, but not limited to, norbornyl, adamantyl, bornyl, camphenyl, isocamphenyl, and carenyl groups. Representative substituted cycloalkyl groups may be mono-substituted or substituted more than once, such as, but not limited to, 2,2-, 2,3-, 2,4- 2,5- or 2,6-disubstituted cyclohexyl groups or mono-, di- or tri-substituted norbornyl or cycloheptyl groups, which may be substituted with, for example, alkyl, alkoxy, amino, thio, cyano, or halo groups.

[0071] Alkenyl groups are straight chain, branched or cyclic lower alkyl groups having 2 to about 8 carbon atoms, and further including at least one double bond, as exemplified, for instance, by vinyl, propenyl, 2-butenyl, 3-butenyl, isobutenyl, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl groups among others.

[0072] Alkynyl groups are straight chain or branched lower alkyl groups having 2 to about 8 carbon atoms, and further including at least one triple bond, as exemplified by groups, including, but not limited to, ethynyl, propynyl, and butynyl groups.

[0073] Aryl groups are cyclic aromatic hydrocarbons that do not contain heteroatoms. Thus aryl groups include, but are not limited to, phenyl, azulene, heptalene, biphenylene, indacene, fluorene, phenanthrene, triphenylene, pyrene, naphthacene, chrysene, biphenyl, anthracenyl, and naphthenyl groups. Although the phrase "aryl groups" includes groups containing fused rings, such as fused aromatic-aliphatic ring systems, it does not include aryl groups that have other groups, such as alkyl or halo groups, bonded to one of the ring members. Rather, groups such as tolyl are referred to as substituted aryl groups. Representative substituted aryl groups may be mono-substituted or substituted more than once, such as, but not limited to, 2-, 3-, 4-, 5-, or 6-substituted phenyl or benzyl groups, which may be substituted with groups including, but not limited to, amino, alkoxy, alkyl, cyano, or halo.

[0074] Cycloalkylalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to a cycloalkyl group as defined above.

[0075] Arylalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above.

[0076] Heterocyclyl groups are nonaromatic ring compounds containing 3 or more ring members, of which, one or more is a heteroatom such as, but not limited to, N, O, and S. The phrase "heterocyclyl group" includes fused ring species including those comprising fused aromatic and nonaromatic groups. The phrase also includes polycyclic ring systems containing a heteroatom such as, but not limited to, quinuclidyl. However, the phrase does not include heterocyclyl groups that have other groups, such as alkyl or halo

groups, bonded to one of the ring members. Rather, these are referred to as "substituted heterocyclyl groups". Heterocyclyl groups include, but are not limited to, piperazino, morpholino, thiomorpholino, pyrrolidino, piperidino and homopiperazino groups. Representative substituted heterocyclyl groups may be mono-substituted or substituted more than once, such as, but not limited to, morpholino or piperazino groups, which are 2-, 3-, 4-, 5-, or 6-substituted, or disubstituted with groups including, but not limited to, amino, alkoxy, alkyl, cyano, or halo.

[0077] Heteroaryl groups are aromatic ring compounds containing 3 or more ring members, of which, one or more is a heteroatom such as, but not limited to, N, O, and S. Heteroaryl groups include, but are not limited to, groups such as furan, thiophene, pyrrole, isopyrrole, diazole, imidazole, isoimidazole, triazole, dithiole, oxathiole, isoxazole, oxazole, thiazole, isothiazole, oxadiazole, oxatriazole, dioxazole, oxathiazole, pyran, dioxin, pyridine, pyrimidine, pyridazine, pyrazine, triazine, oxazine, isoxazine, oxathiazine, azepin, oxepin, thiepin, diazepine, benzofuran, and isobenzofuran. Although the phrase "heteroaryl groups" includes fused ring compounds, the phrase does not include heteroaryl groups that have other groups bonded to one of the ring members, such as alkyl groups. Rather, heteroaryl groups with such substitution are referred to as "substituted heteroaryl groups". Representative substituted heteroaryl groups may be substituted one or more times with groups including, but not limited to, amino, alkoxy, alkyl, cyano, or halo.

[0078] Heterocyclylalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to a heterocyclyl group as defined above.

[0079] Heteroarylalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to a heteroaryl group as defined above.

[0080] Aminocarbonyl groups are groups of the formula $RR'NC(O)-$, wherein R or R' may be the same or different, and each is independently selected from H, or substituted or unsubstituted alkyl, cycloalkyl, aryl, heterocyclyl or heteroaryl groups, as defined above.

[0081] In general, "substituted" refers to a group as defined above in which one or more bonds to a hydrogen atom contained therein are replaced by a bond to non-hydrogen or non-carbon atoms such as, but not limited to, a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, and ester groups; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylarylamines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as in trialkylsilyl groups, dialkylarylsilyl groups, alkylidiarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. Substituted alkyl groups and also substituted cycloalkyl groups and others also include groups in which one or more bonds to a carbon(s) or hydrogen(s) atom is replaced by a bond to a heteroatom such as oxygen in carbonyl, carboxyl, and ester groups; nitrogen in groups such as imines, oximes, hydrazones, and nitriles.

[0082] Substituted cycloalkyl, substituted aryl, substituted heterocyclyl and substituted heteroaryl also include rings and fused ring systems in which a bond to a hydrogen atom is replaced with a bond to a carbon atom. Therefore, substituted cycloalkyl, substituted aryl, substituted heterocyclyl and substituted heteroaryl groups may also be substituted with alkyl groups as defined above.

[0083] Pharmaceutically acceptable salts include a salt with an inorganic base, organic base, inorganic acid, organic acid, or basic or acidic amino acid. As salts of inorganic bases, the invention includes, for example, alkali metals such as sodium or potassium, alkali earth metals such as

calcium and magnesium or aluminum, and ammonia. As salts of organic bases, the invention includes, for example, trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine. As salts of inorganic acids, the instant invention includes, for example, hydrochloric acid, hydroboric acid, nitric acid, sulfuric acid, and phosphoric acid. As salts of organic acids, the instant invention includes, for example, formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, and p-toluenesulfonic acid. As salts of basic amino acids, the instant invention includes, for example, arginine, lysine and ornithine. Acidic amino acids include, for example, aspartic acid and glutamic acid.


[0084] The term "protected" with respect to hydroxyl groups, amine groups, and sulfhydryl groups refers to forms of these functionalities which are protected from undesirable reaction with a protecting group known to those skilled in the art such as those set forth in Protective Groups in Organic Synthesis, Greene, T.W.; Wuts, P. G. M., John Wiley & Sons, New York, NY, (3rd Edition, 1999) which can be added or removed using the procedures set forth therein. Examples of protected hydroxyl groups include, but are not limited to, silyl ethers such as those obtained by reaction of a hydroxyl group with a reagent such as, but not limited to, t-butyldimethyl-chlorosilane, trimethylchlorosilane, triisopropylchlorosilane, triethylchlorosilane; substituted methyl and ethyl ethers such as, but not limited to methoxymethyl ether, methylthiomethyl ether, benzyloxymethyl ether, t-butoxymethyl ether, 2-methoxyethoxymethyl ether, tetrahydropyranyl ethers, 1-ethoxyethyl ether, allyl ether, benzyl ether; esters such as, but not limited to, benzoylformate, formate, acetate, trichloroacetate, and trifluoroacetate. Examples of protected amine groups include, but are not limited to, amides such as, formamide, acetamide, trifluoroacetamide, and benzamide; imides, such as phthalimide, and dithiosuccinimide; and others. Examples of protected sulfhydryl groups include, but are not limited to, thioethers such as S-benzyl thioether, and S-4-

picolyl thioether; substituted S-methyl derivatives such as hemithio, dithio and aminothio acetals; and others.


[0085] Prodrugs, as used in the context of the instant invention, includes those derivatives of the instant compounds which undergo in vivo metabolic biotransformation, by enzymatic or nonenzymatic processes, such as hydrolysis, to form a compound of the invention. Prodrugs can be employed to improve pharmaceutical or biological properties, as for example solubility, melting point, stability and related physicochemical properties, absorption, pharmacodynamics and other delivery-related properties.

[0086] The instant invention provides potent and specific agonists of MC4-R that are low molecular weight small molecules. In accordance with one aspect of the invention, the invention provides compounds of formula IA, IB, and IC. Compounds of the invention further include prodrugs of compounds of formula IA, IB, and IC, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.


[0087] Compounds of formula IA, IB, and IC have the following structure.



IA



IB



IC

[0088] In compounds of formula IA, IB, and IC, Z^1 is selected from the group consisting of CR^4 and N. In some embodiments of the compounds of formula IA, IB, and IC, Z^1 is a CR^4 group whereas in other embodiments, Z^1 is a N.

[0089] In compounds of formula IA, IB, and IC, Z^2 is selected from the group consisting of CR^5 and N. In some embodiments of the compounds of formula IA, IB, and IC, Z^2 is a CR^5 group whereas in other embodiments, Z^2 is a N.

[0090] In compounds of formula IA, IB, and IC, Z^3 is selected from the group consisting of CR^6 and N. In some embodiments of the compounds of formula IA, IB, and IC, Z^3 is a CR^6 group whereas in other embodiments, Z^3 is a N.

[0091] In some embodiments of compounds of formula IA, IB, and IC, Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group. Thus, in some embodiments of compounds of formula IA, IB, and IC, the ring that includes Z^1 , Z^2 , and Z^3 may be a carbocyclic aromatic ring. In some embodiments of compounds of formula IA, IB, and IC, where Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group, at least one of R^4 , R^5 , or R^6 is a halogen such as Cl or F. In other such embodiments, at least one of R^4 , R^5 , or R^6 is a F.

[0092] In some embodiments of compounds of formula IA, IB, and IC, at least one of Z^1 , Z^2 , or Z^3 is a N.

[0093] In compounds of formula IA, IB, and IC, R^1 is selected from the group consisting of substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups. In some embodiments of the compounds of formula IA, IB, and IC, R^1 is a 2,4-disubstituted phenylethyl group. In other embodiments of the compounds of formula IA, IB, and IC, R^1 is selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still other embodiments of the compounds of formula IA, IB, and IC, R^1 is selected from the group that includes phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted

(aryl)(aryloxymethyl)ethyl, substituted and unsubstituted
(aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
(aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted
(aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted
(heteroaryl)(heterocyclyloxymethyl)ethyl groups.

[0094] In some embodiments, the invention provides compounds of formula IA, IB and IC in which R^1 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.


[0095] In compounds of formula IB, R^2 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, alkylcarbonyl, and arylcarbonyl groups. In some embodiments of the compounds of formula IB, R^2 is H.

[0096] In compounds of formula IA and IC, R^3 is selected from the group consisting of H, and substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups. In some embodiments of the compounds of formula IA and IC, R^3 is H. In still further embodiments, the invention provides compounds of formula IA and IC in which R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in which R^3 is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R^3


is selected from substituted and unsubstituted 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0097] In compounds of formula IA, IB, and IC, R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclylamino, heteroaryl amino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl groups. In some embodiments of compounds of formula IA, IB, and IC, R^4 , R^5 , and R^6 are selected from H or a halogen such as Cl or F. In other embodiments of compounds of formula IA, IB, and IC, at least one of R^4 , R^5 , or R^6 is a F whereas in other embodiments of compounds of formula IA, IB, and IC, R^4 , R^5 , and R^6 are all H.

[0098] In compounds of formula IA, IB, and IC, W is a group of formula IIA or IIB having the following structure.



IIA



IIB


[0099] In compounds of formula IA, IB, and IC, $R^{1'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups, and R^2 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In compounds of formula IA, IB, and IC, $R^{1'}$ and R^2 , together with the nitrogen atom to which they are both bound, may alternatively form a substituted or

unsubstituted heterocyclyl or heteroaryl group. In one embodiment of compounds of formula IA, IB, and IC, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still further embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In yet other embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In other embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R^1 and R^2 are bound. In another embodiment of compounds of formula IA, IB, and IC, R^1 and R^2 , together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least two nitrogen atoms. In still another embodiment of compounds of formula IA, IB, and IC, R^1 and R^2 , together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least one oxygen atom and one nitrogen atom. In yet other embodiments of compounds of formula IA, IB, and IC, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino,

homopiperazino, or azepino group. In still further embodiments of compounds of formula IA, IB, and IC, R¹ and R², together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two alkyl groups or in one embodiment by one or two methyl groups.


[0100] In compounds of formula IA, IB, and IC, R³ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In one embodiment of compounds of formula IA, IB, and IC, R³ is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In still other embodiments of compounds of formula IA, IB, and IC, R³ is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkylnorbornyl, bornyl, norbornyl, and decalinyl groups. In still other embodiments of compounds of formula IA, IB, and IC, R³ is selected from the group consisting

of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups. By way of nonlimiting example, suitable R^3 cycloalkyl, cyclohexyl, and polycyclic cycloalkyl groups that include fluorine, include, but are not limited to, the following structures:




[0101] In compounds of formula IA, IB, and IC, R^4 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl groups. In one embodiment of compounds of formula IA, IB, and IC, R^4 is H.


[0102] In one embodiment of the compounds of formula IA, the compounds of formula IA are selected from the group consisting of compounds having the formula IIIA, IIIB, IIIC, IIID, IIIE, IIIF, and IIIG such as shown below where R^1 , W, $R^{1'}$, $R^{2'}$, $R^{3'}$, and $R^{4'}$ have the same values as those described above with respect to the compounds and various embodiments of formula IA, IB, and IC.




IIIA




IIIB




IIIC




IIID



IIIE




IIIF




IIIG

[0103] In an embodiment of the compounds of formula IB, the compounds of formula IB are selected from the group consisting of compounds having the formula IVA and IVB such as shown below where R^1 ,

W, R¹, R², R³, and R⁴ have the same values as those described above with respect to the compounds of formula IA, IB, and IC.




IVA




IVB

[0104] The instant invention provides a second group of compounds that are potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, in accordance with one aspect of the invention, the invention provides a second group of compounds of formula IA and IC. Compounds of the invention further include prodrugs of compounds of formula IA and IC, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0105] Compounds of formula IA and IC have the following structure.



IA



IC

[0106] In the second group of compounds of formula IA and IC, Z¹ is selected from the group consisting of CR⁴ and N. In some embodiments of the second compounds of formula IA and IC, Z¹ is a CR⁴ group whereas in other embodiments, Z¹ is a N.

[0107] In the second group of compounds of formula IA and IC, Z² is selected from the group consisting of CR⁵ and N. In some embodiments of

the second group of compounds of formula IA and IC, Z^2 is a CR^5 group whereas in other embodiments, Z^2 is a N.

[0108] In the second group of compounds of formula IA and IC, Z^3 is selected from the group consisting of CR^6 and N. In some embodiments of the second group of compounds of formula IA and IC, Z^3 is a CR^6 group whereas in other embodiments, Z^3 is a N.

[0109] In some embodiments of the second group of compounds of formula IA and IC, Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group. Thus, in some embodiments of the second group of compounds of formula IA and IC, the ring that includes Z^1 , Z^2 , and Z^3 may be a carbocyclic aromatic ring. In some embodiments of the second group of compounds of formula IA and IC, where Z^1 is a CR^4 group, Z^2 is a CR^5 group, and Z^3 is a CR^6 group, at least one of R^4 , R^5 , or R^6 is a halogen such as Cl or F. In other such embodiments, at least one of R^4 , R^5 , or R^6 is a F.

[0110] In some embodiments of the second group of compounds of formula IA and IC, at least one of Z^1 , Z^2 , or Z^3 is a N.

[0111] In the second group of compounds of formula IA and IC, R^1 is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups. In some embodiments of the second group of compounds of formula IA and IC, R^1 is a 2,4-disubstituted phenylethyl group. In other embodiments of the second group of compounds of formula IA and IC, R^1 is selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^1 is selected from the group that includes phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-

methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups. In some embodiments of the second group of compounds, R^1 is a H or is an alkyl group having from one to eight carbon atoms. In some such embodiments, R^1 is H whereas in other such embodiments, R^1 is a methyl, ethyl, or propyl group. In some such embodiments, R^1 is a methyl group.

[0112] In still further embodiments, the invention provides compounds of formula IA and IC in which R^1 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0113] In the second group of compounds of formula IA and IC, R^3 is selected from the group consisting of substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, alkoxyalkyl, aryloxyalkyl, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl,

alkenyl, alkynyl, alkyl groups, and $-C(=NH)$ -heterocyclyl groups, and groups of formula $-LR^7$. In some embodiments of the second group of compounds of formula IA and IC, R^3 is a 2,4-disubstituted phenylethyl group. In other embodiments of the second group of compounds of formula IA and IC, R^3 is selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R^3 is selected from the group that includes phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxyethyl)ethyl, substituted and unsubstituted (aryl)(heterocycloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxyethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocycloxyethyl)ethyl groups. In some embodiments of the second group of compounds, R^3 has any of the values described in this

paragraph, and R^1 is H or is a substituted or unsubstituted alkyl group. In some such embodiments, R^1 is H.

[0114] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R^3 is a substituted or unsubstituted alkyl group such as a substituted or unsubstituted aryloxyalkyl group or a substituted or unsubstituted heteroaryloxyalkyl group. In some such embodiments, R^3 is a substituted or unsubstituted aryloxymethyl group. In some such embodiments, of the second group of compounds of formula IA and IC, R^3 is selected from the group consisting of $-CH_2-O$ -aryl groups where the aryl group is substituted with one or more halogen group such as with one or more Cl or F. In some such embodiments, the aryl group is additionally substituted with an alkoxy group such as a methoxy or ethoxy group. In some embodiments of the second group of compounds of formula IA and IC, R^3 is a $-CH_2-O$ -aryl group where the aryl group is selected from the group consisting of 2,4-difluorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-methoxyphenyl, 2,4-dichlorophenyl, 4-chlorophenyl, 2-chlorophenyl, and 2-chloro-4-methoxyphenyl groups. In other embodiments of the second group of compounds of formula IA and IC, R^3 is a heterocyclalkyl group. In some other embodiments, of compounds of formula IA and IC, R^3 is a substituted or unsubstituted arylalkoxyalkyl group or a heteroarylalkoxyalkyl group.

[0115] In some aspects of the invention of compounds of the second group of potent and specific agonists of MC4-R, the invention further provides compounds of formula IA and IC in which R^3 is a substituted or unsubstituted heterocyclalkyl group. In some such embodiments, R^3 is a substituted or unsubstituted heterocyclmethyl group. In some such embodiments, the heterocycl group is selected from the group consisting of substituted and unsubstituted 1*H*-tetrazole, piperazine, piperidine, imidazole, and morpholine groups. In some such embodiments, R^3 is a $-CH_2$ -heterocycl group where the heterocycl group is a 1*H*-tetrazole, an imidazole, an N-methylpiperazine,

a 4-hydroxypiperidine, a 3-hydroxypiperidine, or a morpholine. In still other embodiments of the second group of compounds of formula IA and IC, R^3 is a heterocyclyl group. In some embodiments of the second group of compounds of formula IA and IC, R^3 is a substituted or unsubstituted piperazinyl group such as an N-methylpiperazinyl group, is a substituted or unsubstituted pyridine group, is a substituted or unsubstituted tetrazole group, is a substituted or unsubstituted cycloalkyl group such as a 4-methylcyclohexyl group, or is a substituted or unsubstituted phenyl group. In yet other embodiments of the second group of compounds of formula IA and IC, R^3 is an alkoxyalkyl group such as a methoxyalkyl group or an ethoxyalkyl group. In some such embodiments, R^3 is an alkoxyalkyl group such as an alkoxymethyl group such as a methoxymethyl group.


[0116] In some other embodiments of the second group of compounds of formula IA and IC, R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in which R^3 is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R^3 is selected from substituted and unsubstituted 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0117] In some embodiments of the second group of compounds of formula IA and IC, R^1 is an arylalkyl group such as those described above for R^3 . In some such embodiments, R^1 is a substituted or unsubstituted phenylethyl group and R^3 is an alkyl group such as a methyl group. In other embodiments of the second group of compounds, R^1 is an alkyl group such as a methyl group and R^3 is selected from substituted aryloxyalkyl groups, phenylaminoalkyl groups or groups of $-LR^7$ where R^7 is a group formula IIC.


[0118] In the second group of compounds of formula IA and IC, R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br,

OH, NH₂, CN, NO₂, and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclamino, heteroarylmino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclaminocarbonyl, and heteroarylaminocarbonyl groups. In some embodiments of the second group of compounds of formula IA and IC, R⁴, R⁵, and R⁶ are selected from H or a halogen such as Cl or F. In other embodiments of the second group of compounds of formula IA and IC, at least one of R⁴, R⁵, or R⁶ is a F whereas in other embodiments of the second group of compounds of formula IA and IC, R⁴, R⁵, and R⁶ are all H.

[0119] In the second group of compounds of formula IA and IC, W is a group of formula IIA or IIB having the following structure.



IIA



IIB

[0120] In the second group of compounds of formula IA and IC, R^{1'} is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups, and R^{2'} is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In the second group of compounds of formula IA and IC, R^{1'} and R^{2'}, together with the nitrogen atom to which they are both bound, may alternatively form a substituted or unsubstituted heterocyclyl or heteroaryl group. In one embodiment of the second group of compounds of formula IA and IC, R^{1'} is H and R^{2'} is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other

embodiments of the second group of compounds of formula IA and IC, $R^{1'}$ is H and $R^{2'}$ is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still further embodiments of second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$ may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In yet other embodiments of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$ may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In other embodiments of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which $R^{1'}$ and $R^{2'}$ are bound. In another embodiment of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$, together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least two nitrogen atoms. In still another embodiment of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$, together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least one oxygen atom and one nitrogen atom. In yet other embodiments of the second group of compounds of formula IA and IC, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still further embodiments of the second group of compounds of formula IA and IC,

R¹ and R², together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two alkyl groups or in one embodiment by one or two methyl groups.


[0121] In the second group of compounds of formula IA and IC, R³ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In one embodiment of the second group of compounds of formula IA, and IC, R³ is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In still other embodiments of the second group of compounds of formula IA and IC, R³ is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl, bornyl, norbornyl, and decalyl groups. In still other embodiments of the second group of compounds of formula IA and IC, R³ is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-

methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups. By way of nonlimiting example, suitable R³ cycloalkyl, cyclohexyl, and polycyclic cycloalkyl groups that include fluorine, include, but are not limited to, the structures set forth above with respect to the first group of compounds of formula IA, IB, and IC.

[0122] In the second group of compounds of formula IA and IC, R⁴ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocycl, arylalkyl, and heteroarylalkyl groups. In one embodiment of the second group of compounds of formula IA and IC, R⁴ is H.

[0123] In the second group of compounds of formula IA and IC, L is selected from the group consisting of a covalent bond, -CH₂-, -O-, -S-, and -NH-.

[0124] In the second group of compounds of formula IA and IC, R⁷ is selected from the group consisting of substituted and unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from a group of formula IIC;



IIC


[0125] In the second group of compounds of formula IA and IC, Y is selected from the group consisting of CH₂, O, S, and NR⁹ where R⁹. In some embodiments, Y is an NR⁹ group and in some such embodiments, m is 1. In some such embodiments, R⁹ is an alkyl group such as a methyl group or is a H.

[0126] In the second group of compounds of formula IA and IC, R⁸ is selected from the group consisting of H, a halogen, hydroxyl, carboxylic acid, and substituted or unsubstituted alkyl, amino, alkylamino, dialkylamino, alkylaminoalkyl, heterocyclyl, alkoxy, carbonyl, and aminocarbonyl groups.

[0127] In the second group of compounds of formula IA and IC, m is an integer selected from the group consisting of 0, 1, and 2. In some embodiments, m is 1.

[0128] The instant invention provides a third group of compounds that are potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, in accordance with one aspect of the invention, the invention provides a third group of compounds of formula ID. Compounds of the invention further include prodrugs of compounds of formula ID, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0129] Compounds of formula ID have the following structure:



In some embodiments of the third group of compounds of formula ID, at least one of R⁴, R⁵, or R⁶ is a halogen such as Cl or F. In other such embodiments, at least one of R⁴, R⁵, or R⁶ is a F.

[0130] In the third group of compounds of formula ID, R¹ is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups. In some embodiments of the third group of compounds of formula ID, R¹ is a 2,4-disubstituted phenylethyl group. In other embodiments of the third group of compounds of formula ID, R¹ is selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still other embodiments of the third group of compounds of formula ID, R¹ is selected from the group that includes phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted

(heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
(aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
(aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
(aryl)(aryloxymethyl)ethyl, substituted and unsubstituted
(aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
(aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted
(aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
(heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted
(heteroaryl)(heterocyclyloxymethyl)ethyl groups. In some embodiments of the
third group of compounds, R^1 is a H or is an alkyl group having from one to
eight carbon atoms. In some such embodiments, R^1 is H whereas in other
such embodiments, R^1 is a methyl, ethyl, or propyl group. In some such
embodiments, R^1 is a methyl group.

[0131] In still further embodiments, the invention provides compounds
of formula ID in which R^1 is a substituted or unsubstituted alkenyl group such
as a substituted or unsubstituted allyl group or a substituted or unsubstituted
vinyl group.

[0132] In the third group of compounds of formula ID, R^2 and R^3 are
independently selected from the group consisting of substituted and
unsubstituted arylalkyl, heteroarylalkyl, alkoxy, alkoxyalkyl, aryloxyalkyl,
alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl,
heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkyl groups, and $-C(=NH)-$
heterocyclyl groups, and groups of formula $-LR^7$. In some embodiments of the
third group of compounds of formula ID, R^2 is -H or a substituted or
unsubstituted alkyl group. In some embodiments of the third group of
compounds of formula ID, R^3 is a 2,4-disubstituted phenylethyl group. In
other embodiments of the third group of compounds of formula ID, R^3 is
selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still

other embodiments of the third group of compounds of formula ID, R^3 is selected from the group that includes phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocyclyloxymethyl)ethyl groups. In some embodiments of the third group of compounds, R^3 has any of the values described in this paragraph, and R^1 or R^2 is H or is a substituted or unsubstituted alkyl group. In some such embodiments, R^1 is H. In other such embodiments, R^2 is H.

[0133] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^3 is a substituted or unsubstituted alkyl group such as a substituted or unsubstituted aryloxyalkyl group or a

substituted or unsubstituted heteroaryloxyalkyl group. In some such embodiments, R^3 is a substituted or unsubstituted aryloxymethyl group. In some such embodiments, of the third group of compounds of formula ID, R^3 is selected from the group consisting of $-CH_2-O$ -aryl groups where the aryl group is substituted with one or more halogen group such as with one or more Cl or F. In some such embodiments, the aryl group is additionally substituted with an alkoxy group such as a methoxy or ethoxy group. In some embodiments of the third group of compounds of formula ID, R^3 is a $-CH_2-O$ -aryl group where the aryl group is selected from the group consisting of 2,4-difluorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-methoxyphenyl, 2,4-dichlorophenyl, 4-chlorophenyl, 2-chlorophenyl, and 2-chloro-4-methoxyphenyl groups. In other embodiments of the third group of compounds of formula ID, R^3 is a heterocyclylalkyl group. In some other embodiments, of compounds of formula ID, R^3 is a substituted or unsubstituted arylalkoxyalkyl group or a heteroarylalkoxyalkyl group.

[0134] In some aspects of the invention of compounds of the third group of potent and specific agonists of MC4-R, the invention further provides compounds of formula ID in which R^3 is a substituted or unsubstituted heterocyclylalkyl group. In some such embodiments, R^3 is a substituted or unsubstituted heterocyclylmethyl group. In some such embodiments, the heterocyclyl group is selected from the group consisting of substituted and unsubstituted 1*H*-tetrazole, piperazine, piperidine, imidazole, and morpholine groups. In some such embodiments, R^3 is a $-CH_2$ -heterocyclyl group where the heterocyclyl group is a 1*H*-tetrazole, an imidazole, an N-methylpiperazine, a 4-hydroxypiperidine, a 3-hydroxypiperidine, or a morpholine. In still other embodiments of the third group of compounds of formula ID, R^3 is a heterocyclyl group. In some embodiments of the third group of compounds of formula ID, R^3 is a substituted or unsubstituted piperazinyl group such as an N-methylpiperazinyl group, is a substituted or unsubstituted pyridine group, is a substituted or unsubstituted tetrazole group, is a substituted or unsubstituted cycloalkyl group such as a 4-methylcyclohexyl group, or is a

substituted or unsubstituted phenyl group. In yet other embodiments of the third group of compounds of formula ID, R^3 is an alkoxyalkyl group such as a methoxyalkyl group or an ethoxyalkyl group. In some such embodiments, R^3 is a an alkoxyalkyl group such as an alkoxymethyl group such as a methoxymethyl group.


[0135] In some other embodiments of the third group of compounds of formula ID, R^3 is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group. In some embodiments, the invention provides compounds in which R^3 is a substituted or unsubstituted heteroaryl group, heterocyclyl group, alkylamino group, or cycloalkyl amino group. In some such embodiments, R^3 is selected from substituted and unsubstituted 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazine, morpholinyl, piperazinyl, and cyclopropylamino groups.

[0136] In some embodiments of the third group of compounds of formula ID, R^1 is an arylalkyl group such as those described above for R^3 . In some such embodiments, R^1 is a substituted or unsubstituted phenylethyl group and R^3 is an alkyl group such as a methyl group. In other embodiments of the third group of compounds, R^1 is an alkyl group such as a methyl group and R^3 is selected from substituted aryloxyalkyl groups, phenylaminoalkyl groups or groups of $-LR^7$ where R^7 is a group formula IIC.

[0137] In the third group of compounds of formula ID, R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclylamino, heteroarylamino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl groups. In some embodiments of the third group of compounds of formula ID, R^4 , R^5 , and R^6 are selected from H or a halogen such as Cl or F. In other embodiments of the third group of compounds of

formula ID, at least one of R^4 , R^5 , or R^6 is a F whereas in other embodiments of the third group of compounds of formula ID, R^4 , R^5 , and R^6 are all H.

[0138] In the third group of compounds of formula ID, W is a group of formula IIA or IIB having the following structure.



[0139] In the third group of compounds of formula ID, $R^{1'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups, and $R^{2'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen atom to which they are both bound, may alternatively form a substituted or unsubstituted heterocyclyl or heteroaryl group. In one embodiment of the third group of compounds of formula ID, $R^{1'}$ is H and $R^{2'}$ is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the third group of compounds of formula ID, $R^{1'}$ is H and $R^{2'}$ is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still further embodiments of third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$ may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In yet other

embodiments of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$ may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In other embodiments of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which $R^{1'}$ and $R^{2'}$ are bound. In another embodiment of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least two nitrogen atoms. In still another embodiment of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least one oxygen atom and one nitrogen atom. In yet other embodiments of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still further embodiments of the third group of compounds of formula ID, $R^{1'}$ and $R^{2'}$, together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two alkyl groups or in one embodiment by one or two methyl groups.

[0140] In the third group of compounds of formula ID, $R^{3'}$ is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In one embodiment of the third group of compounds of formula IA, and IC, $R^{3'}$ is selected from the


group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In still other embodiments of the third group of compounds of formula ID, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl, bornyl, norbornyl, and decalanyl groups. In still other embodiments of the third group of compounds of formula ID, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl)

groups. By way of nonlimiting example, suitable R^3 cycloalkyl, cyclohexyl, and polycyclic cycloalkyl groups that include fluorine, include, but are not limited to, the structures set forth above with respect to the first group of compounds of formula IA, IB, and IC.

[0141] In the third group of compounds of formula ID, $R^{4'}$ is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, aryl, heteroaryl, heterocycl, arylalkyl, and heteroarylalkyl groups. In one embodiment of the third group of compounds of formula ID, $R^{4'}$ is H.

[0142] In the third group of compounds of formula ID, L is selected from the group consisting of a covalent bond, $-CH_2-$, $-O-$, $-S-$, and $-NH-$.

[0143] In the third group of compounds of formula ID, R^7 is selected from the group consisting of substituted and unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from a group of formula IIC;



IIC


[0144] In the third group of compounds of formula ID, Y is selected from the group consisting of CH_2 , O, S, and NR^9 where R^9 . In some embodiments, Y is an NR^9 group and in some such embodiments, m is 1. In some such embodiments, R^9 is an alkyl group such as a methyl group or is a H.

[0145] In the third group of compounds of formula ID, R^8 is selected from the group consisting of H, a halogen, hydroxyl, carboxylic acid, and substituted or unsubstituted alkyl, amino, alkylamino, dialkylamino, alkylaminoalkyl, heterocycl, alkoxy, carbonyl, and aminocarbonyl groups.

[0146] In the third group of compounds of formula ID, m is an integer selected from the group consisting of 0, 1, and 2. In some embodiments, m is 1.

[0147] The instant invention provides a fourth group of compounds that are potent and specific agonists of MC4-R that are low molecular weight small molecules. Thus, in accordance with one aspect of the invention, the invention provides a fourth group of compounds of formula IE. Compounds of the invention further include prodrugs of compounds of formula IE, pharmaceutically acceptable salts thereof, stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof, or solvates thereof.

[0148] Compounds of formula IE have the following structure.



[0149] In some embodiments of the fourth group of compounds of formula IE at least one of R^4 , R^5 , or R^6 is a halogen such as Cl or F. In other such embodiments, at least one of R^4 , R^5 , or R^6 is a F.

[0150] In the fourth group of compounds of formula IE, R^1 is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups. In some embodiments of the fourth group of compounds of formula IE, R^1 is a 2,4-disubstituted phenylethyl group. In other embodiments of the fourth group of compounds of formula IE, R^1 is selected from 2,4-dihalophenylethyl and 2,4-dialkylphenylethyl groups. In still other embodiments of the fourth group of compounds of formula IE, R^1 is selected from the group that includes


phenylethyl, 2,4-dichlorophenylethyl, 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl, (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted (aryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted (aryl)(heteroaryloxyethyl)ethyl, substituted and unsubstituted (aryl)(heterocycloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(alkoxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted (heteroaryl)(heteroaryloxyethyl)ethyl, and substituted and unsubstituted (heteroaryl)(heterocycloxyethyl)ethyl groups. In some embodiments of the fourth group of compounds, R¹ is a H or is an alkyl group having from one to eight carbon atoms. In some such embodiments, R¹ is H whereas in other such embodiments, R¹ is a methyl, ethyl, or propyl group. In some such embodiments, R¹ is a methyl group.

[0151] In still further embodiments, the invention provides compounds of formula IE in which R¹ is a substituted or unsubstituted alkenyl group such as a substituted or unsubstituted allyl group or a substituted or unsubstituted vinyl group.

[0152] In the fourth group of compounds of formula IE, R^1 is H or is a substituted or unsubstituted alkyl group. In some such embodiments, R^1 is H.

[0153] In the fourth group of compounds of formula IE, R^4 , R^5 , and R^6 are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclamino, heteroarylmino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclaminocarbonyl, and heteroarylaminocarbonyl groups. In some embodiments of the fourth group of compounds of formula IE, R^4 , R^5 , and R^6 are selected from H or a halogen such as Cl or F. In other embodiments of the fourth group of compounds of formula IE, at least one of R^4 , R^5 , or R^6 is a F whereas in other embodiments of the fourth group of compounds of formula IE, R^4 , R^5 , and R^6 are all H.

[0154] In the fourth group of compounds of formula IE, W is a group of formula IIA or IIB having the following structure.



[0155] In the fourth group of compounds of formula IE, R^1 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups, and R^2 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen atom to which they are both bound, may

alternatively form a substituted or unsubstituted heterocyclyl or heteroaryl group. In one embodiment of the fourth group of compounds of formula IE, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In still other embodiments of the fourth group of compounds of formula IE, R^1 is H and R^2 is selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still further embodiments of fourth group of compounds of formula IE, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl groups. In yet other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 may be the same or different and are each independently selected from the group consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and thiophene groups. In still other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted heterocyclyl group. In other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a substituted or unsubstituted saturated heterocyclyl group comprising at least one heteroatom selected from the group consisting of O, S, and N, in addition to the nitrogen atom to which R^1 and R^2 are bound. In another embodiment of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least two nitrogen atoms. In still another embodiment of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen atom to which they are bound, form a substituted or unsubstituted heterocyclyl ring containing at least one oxygen atom and one nitrogen atom. In yet other embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the

nitrogen to which they are bound, form a substituted or unsubstituted piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino group. In still further embodiments of the fourth group of compounds of formula IE, R^1 and R^2 , together with the nitrogen to which they are bound, form a piperazino group optionally substituted by one or two alkyl groups or in one embodiment by one or two methyl groups.

[0156] In the fourth group of compounds of formula IE, R^3 is selected from the group consisting of H, and substituted and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups. In one embodiment of the fourth group of compounds of formula IA, and IC, R^3 is selected from the group consisting of substituted and unsubstituted cycloalkyl, polycyclic cycloalkyl, alkenyl, alkyl, and aryl groups. In still other embodiments of the fourth group of compounds of formula IE, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-alkylcyclohexyl, 2,2-dialkylcyclohexyl, 2,3-dialkylcyclohexyl, 2,4-dialkylcyclohexyl, 2,5-dialkylcyclohexyl, 2,6-dialkylcyclohexyl, 3,4-dialkylcyclohexyl, 3-alkylcyclohexyl, 4-alkylcyclohexyl, 3,3,5-trialkylcyclohexyl, cyclohexylmethyl, 2-aminocyclohexyl, 3-aminocyclohexyl, 4-aminocyclohexyl, 2,3-diaminocyclohexyl, 2,4-diaminocyclohexyl, 3,4-diaminocyclohexyl, 2,5-diaminocyclohexyl, 2,6-diaminocyclohexyl, 2,2-diaminocyclohexyl, 2-alkoxycyclohexyl, 3-alkoxycyclohexyl, 4-alkoxycyclohexyl, 2,3-dialkoxycyclohexyl, 2,4-dialkoxycyclohexyl, 3,4-dialkoxycyclohexyl, 2,5-dialkoxycyclohexyl, 2,6-dialkoxycyclohexyl, 2,2-dialkoxycyclohexyl, 2-alkylthiocyclohexyl, 3-alkylthiocyclohexyl, 4-alkylthiocyclohexyl, 2,3-dialkylthiocyclohexyl, 2,4-dialkylthiocyclohexyl, 3,4-dialkylthiocyclohexyl, 2,5-dialkylthiocyclohexyl, 2,6-dialkylthiocyclohexyl, 2,2-dialkylthiocyclohexyl, fluorocycloalkyl, fluoroalkylcycloalkyl, trifluoromethylcycloalkyl, cyclopentyl, cycloheptyl, cyclohexenyl, isopropyl, n-butyl, cyclooctyl, 2-aryl cyclohexyl, 2-phenylcyclohexyl, 2-arylalkylcyclohexyl, 2-benzylcyclohexyl, 4-phenylcyclohexyl, adamantyl, isocamphenyl, carenyl, 7,7-dialkyl norbornyl,

bornyl, norbornyl, and decaliny groups. In still other embodiments of the fourth group of compounds of formula IE, R^3 is selected from the group consisting of substituted and unsubstituted cyclohexyl, 2-methylcyclohexyl, 2,2-dimethylcyclohexyl, 2,3-dimethylcyclohexyl, 2,4-dimethylcyclohexyl, 2,5-dimethylcyclohexyl, 2,6-dimethylcyclohexyl, 3,4-dimethylcyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, cyclohexenyl, 3,3,5-trimethylcyclohexyl, 4-*t*-butylcyclohexyl, cyclohexylmethyl, isopinocampheyl, 7,7-dimethylnorbornyl, 4-isopropylcyclohexyl, 3-methylcycloheptyl groups, 2-fluoro-4-methylcyclohexyl, 4-fluoro-2-methylcyclohexyl, 4,4-difluoro-2-methylcyclohexyl, 4-trifluoromethylcyclohexyl, 2-methyl-4-trifluoromethylcyclohexyl, 2-fluoromethylcyclohexyl, trifluoromethyl(polycyclic cycloalkyl), fluoromethyl(polycyclic cycloalkyl), and fluoro(polycyclic cycloalkyl) groups. By way of nonlimiting example, suitable R^3 cycloalkyl, cyclohexyl, and polycyclic cycloalkyl groups that include fluorine, include, but are not limited to, the structures set forth above with respect to the first group of compounds of formula IA, IB, and IC.

[0157] In the fourth group of compounds of formula IE, R^4 is selected from the group consisting of H, and substituted and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclialkyl, cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl groups. In one embodiment of the fourth group of compounds of formula IE, R^4 is H.

[0158] In some embodiments of any of the compounds of the invention which includes a W group of formula IIA or IIB where R^1 and R^2 join together with the nitrogen to which they are bound to form a heterocyclic group, the heterocyclic group is substituted with a -CN group, an -OH group, a -CF₃ group, a -CH₂F group, a -CHF₂ group, a -CH₂CN group, a -CH₂OH group, a -CH₂O-alkyl group, or a cycloalkyl group such as a cyclopropyl group. In some such compounds, the heterocyclic compound is a piperidine or a piperazine. In some such compounds, R^1 and R^2 join together with the nitrogen to which they are bound to form a piperazine in which the N atom in the piperazine ring which is not part of the guanidine group is substituted with a -C≡N group, an -

OH group, a $-\text{CH}_2\text{CN}$ group, or a cycloalkyl group. In some compounds in which R^1 and R^2 join together with the nitrogen to which they are bound to form a heterocyclic ring, the heterocycle is a bicyclic structure that includes a spirocenter such that the heterocyclic ring is part of a spirocyclic structure. In some compounds in which R^1 and R^2 join together with the nitrogen to which they are bound to form a heterocyclic ring, the heterocyclic ring is substituted such that a ring carbon atom of the heterocyclic ring is a carbonyl carbon or the carbon of the heterocyclic ring is replaced with a sulfur that is bonded to one or more oxygen atoms. For example, in some embodiments R^1 and R^2 join together with the nitrogen to which they are bound to form a piperazine ring in which one of the ring carbon atoms is a carbonyl carbon atom such that the piperazine compound is a lactam which may be further substituted, for example, with an alkyl group such as a methyl group.

[0159] There has also been provided, in accordance with another aspect of the invention, a composition such as a pharmaceutical formulation or medicament comprising a compound according to the instant invention and a pharmaceutically acceptable carrier. The invention further provides the use of the compounds of the invention in preparing a medicament such as a medicament for use in treating an MC4-R mediated disease. In some embodiments, such a disease is obesity or type II diabetes.

[0160] There has also been provided, in accordance with another aspect of the invention, a method of activating MC4-R in a subject, comprising administering to a subject in need thereof an effective amount of a compound or composition of the instant invention. In some embodiments, the compound or composition is administered to the subject intranasally. In some embodiments, the subject is human.

[0161] There has also been provided, in accordance with another aspect of the invention, a method of treating an MC4-R-mediated disease, comprising administering to a subject in need thereof, a compound or composition of the instant invention. In some embodiments, the compound or

composition is administered to the subject intranasally. In some embodiments, the subject is human

[0162] In one embodiment, a disease to be treated by those methods of the instant invention is obesity, or type I or type II diabetes.

[0163] In another embodiment, a condition to be treated by those methods of the instant invention is a condition associated with or a complication arising from obesity or type II diabetes.

[0164] In another embodiment, a condition to be treated by those methods of the instant invention is erectile dysfunction.

[0165] In another embodiment, a disease to be treated by those methods of the instant invention is polycystic ovary disease.

[0166] In another embodiment, a disease to be treated by those methods of the instant invention is Syndrome X.

[0167] The invention also includes tautomers of the instant compounds. The instant invention also includes prodrugs, pharmaceutically acceptable salts, stereoisomers, hydrates, hydrides, and solvates of these tautomers.

[0168] The instant compounds may exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. In some cases, one stereoisomer may be more active and/or may exhibit beneficial effects in comparison to other stereoisomer(s) or when separated from the other stereoisomer(s). However, it is well within the skill of the ordinary artisan to separate, and/or to selectively prepare said stereoisomers. Accordingly, "stereoisomers" of the instant invention necessarily includes mixtures of stereoisomers, individual stereoisomers, or optically active forms.

[0169] The instant invention also provides for compositions which may be prepared by mixing one or more compounds of the instant invention, or

pharmaceutically acceptable salts or tautomers thereof, with pharmaceutically acceptable carriers, excipients, binders, diluents or the like, to treat or ameliorate a variety of disorders. Examples of such disorders include, but are not limited to obesity, erectile disorders, cardiovascular disorders, neuronal injuries or disorders, inflammation, fever, cognitive disorders, sexual behavior disorders. A therapeutically effective dose further refers to that amount of one or more compounds of the instant invention sufficient to result in amelioration of symptoms of the disorder. The pharmaceutical compositions of the instant invention can be manufactured by methods well known in the art such as conventional granulating, mixing, dissolving, encapsulating, lyophilizing, emulsifying or levigating processes, among others. The compositions can be in the form of, for example, granules, powders, tablets, capsules, syrup, suppositories, injections, emulsions, elixirs, suspensions or solutions. The instant compositions can be formulated for various routes of administration, for example, by oral administration, by intranasal administration, by transmucosal administration, by rectal administration, or subcutaneous administration as well as intrathecal, intravenous, intramuscular, intraperitoneal, intranasal, intraocular or intraventricular injection. The compound or compounds of the instant invention can also be administered in a local rather than a systemic fashion, such as injection as a sustained release formulation. The following dosage forms are given by way of example and should not be construed as limiting the instant invention.

[0170] For oral, buccal, and sublingual administration, powders, suspensions, granules, tablets, pills, capsules, gelcaps, and caplets are acceptable as solid dosage forms. These can be prepared, for example, by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers thereof, with at least one additive or excipient such as a starch or other additive. Suitable additives or excipients are sucrose, lactose, cellulose sugar, mannitol, maltitol, dextran, sorbitol, starch, agar, alginates, chitins, chitosans, pectins, tragacanth gum, gum arabic, gelatins, collagens, casein, albumin, synthetic or semi-synthetic polymers or

glycerides, methyl cellulose, hydroxypropylmethyl-cellulose, and/or polyvinylpyrrolidone. Optionally, oral dosage forms can contain other ingredients to aid in administration, such as an inactive diluent, or lubricants such as magnesium stearate, or preservatives such as parabens or sorbic acid, or anti-oxidants such as ascorbic acid, tocopherol or cysteine, a disintegrating agent, binders, thickeners, buffers, sweeteners, flavoring agents or perfuming agents. Additionally, dyestuffs or pigments may be added for identification. Tablets and pills may be further treated with suitable coating materials known in the art.

[0171] Liquid dosage forms for oral administration may be in the form of pharmaceutically acceptable emulsions, syrups, elixirs, suspensions, slurries and solutions, which may contain an inactive diluent, such as water.

Pharmaceutical formulations may be prepared as liquid suspensions or solutions using a sterile liquid, such as, but not limited to, an oil, water, an alcohol, and combinations of these. Pharmaceutically suitable surfactants, suspending agents, emulsifying agents, may be added for oral or parenteral administration.

[0172] As noted above, suspensions may include oils. Such oils include, but are not limited to, peanut oil, sesame oil, cottonseed oil, corn oil and olive oil. Suspension preparation may also contain esters of fatty acids such as ethyl oleate, isopropyl myristate, fatty acid glycerides and acetylated fatty acid glycerides. Suspension formulations may include alcohols, such as, but not limited to, ethanol, isopropyl alcohol, hexadecyl alcohol, glycerol and propylene glycol. Ethers, such as but not limited to, poly(ethyleneglycol), petroleum hydrocarbons such as mineral oil and petrolatum; and water may also be used in suspension formulations.

[0173] For intranasal administration (e.g., to deliver compounds to the brain), or administration by inhalation (e.g., to deliver compounds through the lungs), the pharmaceutical formulations may be a solution, a spray, a dry powder, or aerosol containing any appropriate solvents and optionally other

compounds such as, but not limited to, stabilizers, antimicrobial agents, antioxidants, pH modifiers, surfactants, bioavailability modifiers and combinations of these. Examples of intranasal formulations and methods of administration can be found in WO 01/41782, WO 00/33813, WO 91/97947, U.S. Patent No. 6,180,603, and U.S. Patent No. 5,624,898. A propellant for an aerosol formulation may include compressed air, nitrogen, carbon dioxide, or a hydrocarbon based low boiling solvent. The compound or compounds of the instant invention are conveniently delivered in the form of an aerosol spray presentation from a nebulizer or the like.

[0174] Injectable dosage forms generally include aqueous suspensions or oil suspensions which may be prepared using a suitable dispersant or wetting agent and a suspending agent. Injectable forms may be in solution phase or in the form of a suspension, which is prepared with a solvent or diluent. Acceptable solvents or vehicles include sterilized water, Ringer's solution, or an isotonic aqueous saline solution. Alternatively, sterile oils may be employed as solvents or suspending agents. Preferably, the oil or fatty acid is non-volatile, including natural or synthetic oils, fatty acids, mono-, di- or tri-glycerides.

[0175] For injection, the pharmaceutical formulation may be a powder suitable for reconstitution with an appropriate solution as described above. Examples of these include, but are not limited to, freeze dried, rotary dried or spray dried powders, amorphous powders, granules, precipitates, or particulates. For injection, the formulations may optionally contain stabilizers, pH modifiers, surfactants, bioavailability modifiers and combinations of these. The compounds may be formulated for parenteral administration by injection such as by bolus injection or continuous infusion. A unit dosage form for injection may be in ampoules or in multi-dose containers.

[0176] For rectal administration, the pharmaceutical formulations may be in the form of a suppository, an ointment, an enema, a tablet or a cream for release of compound in the intestines, sigmoid flexure and/or rectum. Rectal

suppositories are prepared by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers of the compound, with acceptable vehicles, for example, cocoa butter or polyethylene glycol, which is present in a solid phase at normal storing temperatures, and present in a liquid phase at those temperatures suitable to release a drug inside the body, such as in the rectum. Oils may also be employed in the preparation of formulations of the soft gelatin type and suppositories. Water, saline, aqueous dextrose and related sugar solutions, and glycerols may be employed in the preparation of suspension formulations which may also contain suspending agents such as pectins, carbomers, methyl cellulose, hydroxypropyl cellulose or carboxymethyl cellulose, as well as buffers and preservatives.

[0177] Besides those representative dosage forms described above, pharmaceutically acceptable excipients and carriers are generally known to those skilled in the art and are thus included in the instant invention. Such excipients and carriers are described, for example, in "Remingtons Pharmaceutical Sciences" Mack Pub. Co., New Jersey (1991), which is incorporated herein by reference.

[0178] The formulations of the invention may be designed for to be short-acting, fast-releasing, long-acting, and sustained-releasing as described below. Thus, the pharmaceutical formulations may also be formulated for controlled release or for slow release.

[0179] The instant compositions may also comprise, for example, micelles or liposomes, or some other encapsulated form, or may be administered in an extended release form to provide a prolonged storage and/or delivery effect. Therefore, the pharmaceutical formulations may be compressed into pellets or cylinders and implanted intramuscularly or subcutaneously as depot injections or as implants such as stents. Such implants may employ known inert materials such as silicones and biodegradable polymers.

[0180] A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms. Specific dosages may be adjusted depending on conditions of disease, the age, body weight, general health conditions, sex, diet of the subject, dose intervals, administration routes, excretion rate, and combinations of drugs. Any of the above dosage forms containing effective amounts are well within the bounds of routine experimentation and therefore, well within the scope of the instant invention. A therapeutically effective dose may vary depending upon the route of administration and dosage form. The preferred compound or compounds of the instant invention is a formulation that exhibits a high therapeutic index. The therapeutic index is the dose ratio between toxic and therapeutic effects which can be expressed as the ratio between LD₅₀ and ED₅₀. The LD₅₀ is the dose lethal to 50% of the population and the ED₅₀ is the dose therapeutically effective in 50% of the population. The LD₅₀ and ED₅₀ are determined by standard pharmaceutical procedures in animal cell cultures or experimental animals.

[0181] The present invention also provides methods of enhancing MC4-R activity in a human or non-human animal. The method comprises administering an effective amount of a compound, or composition, of the instant invention to said mammal or non-human animal. Effective amounts of the compounds of the instant invention include those amounts that activate MC4-R which are detectable, for example, by an assay described below in the illustrative Examples, or any other assay known by those skilled in the art that a detect signal transduction, in a biochemical pathway, through activation of G-protein coupled receptors, for example, by measuring an elevated cAMP level as compared to a control model. Accordingly, "activating" means the ability of a compound to initiate a detectable signal. Effective amounts may also include those amounts which alleviate symptoms of a MC4-R disorder treatable by activating MC4-R.


[0182] An MC4-R disorder, or MC4-R-mediated disease, which may be treated by those methods provided, include any biological disorder or disease

in which MC4-R is implicated, or which inhibition of MC4-R potentiates a biochemical pathway that is defective in the disorder or disease state. Examples of such diseases are obesity, erectile disorders, cardiovascular disorders, neuronal injuries or disorders, inflammation, fever, cognitive disorders, type II diabetes, polycystic ovary disease, Syndrome X, complications from obesity and diabetes, and sexual behavior disorders. In a preferred embodiment, the instant invention provides compounds, compositions, and methods effective for reducing energy intake and body weight; reducing serum insulin and glucose levels; alleviating insulin resistance; and reducing serum levels of free fatty acids. Accordingly, the instant invention is particularly effective in treating those disorders or diseases associated with obesity or type II diabetes.

[0183] "Treating" within the context of the instant invention, therefore, means an alleviation of symptoms associated with a disorder or disease, or halt of further progression or worsening of those symptoms, or prevention or prophylaxis of the disease or disorder. For example, within the context of obesity, successful treatment may include an alleviation of symptoms or halting the progression of the disease, as measured by reduction in body weight, or a reduction in amount of food or energy intake. In this same vein, successful treatment of type I or type II diabetes may include an alleviation of symptoms or halting the progression of the disease, as measured by a decrease in serum glucose or insulin levels in, for example, hyperinsulinemic or hyperglycemic patients.

[0184] Compounds of formula IA may be readily synthesized as shown in Scheme 1a, the specifics of which are provided in the Examples section. As also explained in the Examples section, compounds of formula IIIC, IIID, and IIIE may be prepared using the methodology shown in Scheme 1b using an appropriately fluorine-substituted 4-nitroanthranilic acid in place of compound (b). Compounds of formula IC may be prepared using the methodology shown in Scheme 1a using an appropriately substituted 5-nitroanthranilic acid in place of 4-nitroanthranilic acid.


SCHEME 1a



[0185] As explained in the Examples section, compounds of formula IIIA, IIIB, IIIF, and IIIG may be prepared using the methodology shown in Scheme 1b using an appropriately substituted pyridine, pyrazine, or pyrimidine carboxylic acid, for example 2-amino-6-azido-nicotinic acid, shown in (b). The conversion of compound (d) to (e), described in further detail in Procedure 1A, proceeds through the initial addition of trimethylphosphine to form a reactive iminophosphorane intermediate, followed by the addition of a substituted isocyanate (preferably by a substituted carbocycle isocyanate) to


produce a carbodiimide, and finally formation of (e) occurs upon reaction with an amine (preferably a substituted piperazine).

Scheme 1b




[0186] Compounds of formula IA may also be prepared using the general synthetic methodology set forth in Scheme 2a. This methodology is also suitable for preparing compounds of formula IC by using a benzoic acid with the nitro group in the correct position.

SCHEME 2a




[0187] Scheme 2b shows an alternative route that may be used to prepare various compounds of formula IA.

SCHEME 2b




[0188] Scheme 2c shows an alternative route that may be used to prepare various compounds of formula IA. This route may also be used to prepare compound of formula IC if a starting material is used with a nitro group in the appropriate position as will be readily observed by those skilled in the art.

SCHEME 2c



[0189] The invention provides various methods for synthesizing compounds of formula IA and IC, various intermediate compounds, and salts of the compounds and intermediate compounds. For example, a method for producing a compound having the formula VA is provided where R¹, R³, Z¹, Z², Z³, R⁴, R⁵, and R⁶ have any of the values described above with respect to compounds of formula IA and embodiments of compounds of formula IA, and Y¹ is selected from the group consisting of NO₂, a protected amine group, a halogen such as Cl, F, Br, or I, an -NCS, and an N₃ group. In some embodiments of the method, R¹ is a substituted or unsubstituted arylalkyl group. In some embodiments of the method, R³ is H. In some embodiments, Y¹ is selected from a F, NO₂, or an N₃ group. In some embodiments of the method, R¹ is a substituted or unsubstituted arylalkyl group, R³ is H, and Y¹ is selected from a F, NO₂, or an N₃ group. In some such embodiments, Z¹, Z²,


and Z^3 are each carbon atoms and R^4 , R^5 , and R^6 are each H. Compounds of formula VA may be readily converted into compounds of formula IA as shown in Scheme 1a and the procedures set forth herein. Typically a compound of formula VA where Y^1 is a $-NCS$ is reacted with a first amine and then is reacted with a second amine as described in Procedure 1a. Any of the amines defined by the R^1 , R^2 , and R^3 of compounds of formula IA may be used. The versatility of this procedure allows a wide range of compounds of formula IA to be prepared where W is a guanidine group prepared from compounds of formula VA where Y^1 is a $-NCS$ group. Another procedure that may be used to prepare compounds of formula IA from compounds of formula VA where Y^1 is an N_3 group is shown in Scheme 1b where nitrogen compounds with any of the R^1 , R^2 , and R^3 groups of compounds of formula IA may be used.



VA


[0190] A method for preparing a compound of formula VA typically includes cyclizing a compound of formula VB by reacting it with an electrophilic carbon bearing an R^3 group such as a trialkyl orthoformate such as triethyl orthoformate, trimethyl orthoformate or the like where the R^3 is a H; Gold's Reagent; a substituted or unsubstituted alkanoyl halide such as acetyl chloride where R^3 is a methyl group; a substituted or unsubstituted alkanoyl acid in the presence of an acid halide producing agent such as thionyl chloride, $POCl_3$, various phosphorous halides, and the like (e.g. an alkanoyl acid of formula R^3CO_2H in combination with $POCl_3$); a benzoyl chloride or an analogous heteroaryl acid chloride compound; or a substituted or unsubstituted benzoic acid or analogous heteroaryl carboxylic acid compound

and an acid halide producing agent. The reaction provides the compound of formula VA. Compounds of formula VB have the following formula.



VB

In compounds of formula VB, the variables may have any of the same values as described above with respect to compounds of formula VA. The method of forming a compound of formula VA may include reacting a compound of formula VC with an amine of formula R^1-NH_2 to produce the compound of formula VB using standard amide-forming procedures and where R^1 has any of the values of the compounds of formula VA and VB and the variables in the compound of VC have the values of compounds of formula VA and VB and Y^2 is a hydroxyl group or is an equivalent thereof. Compounds of formula VC have the following formula.



VC

[0191] An alternative method for synthesizing compounds of formula IA is shown in Scheme 2a. Such a method generally includes reacting a compound of formula VC with an amine of formula R^1-NH_2 and an electrophilic carbon bearing an R^3 group such as a carboxylic acid of formula R^3-CO_2H .


Compounds of formula IA may further be prepared according to the procedure in Scheme 2b by reacting a compound of formula VC with an amide of formula $R^1-N(H)-C(=O)-R^3$ in the presence of $POCl_3$ or an analogous compound.

[0192] Compounds of formula IE may be prepared from compounds of formula VB where Y^1 is a NO_2 group by reacting the compound with $NaNO_2$ as described in Method 6 to produce the analogs of compounds of formula VA which may then be converted to the compounds of formula IE from the compounds where Y^1 is an $-N_3$ group or is an $-NCS$ group.

[0193] Compounds of formula IB may be produced from compounds of formula VB by reacting the compound of formula VB with phosgene or an equivalent thereof as described in Step 2 of Example 2 and subsequent conversion to the guanidine compounds from the N_3 or $-NCS$ compounds using the standard procedures. Finally, compounds of formula ID may be prepared using the procedures described in Method 7 (Steps 1 and 2) using the fragments with the variables described above with respect to compounds of formula ID.

[0194] As noted above, the invention also provides methods for synthesizing compounds of formula IC, various intermediate compounds, and salts of the compounds and intermediate compounds. For example, a method for producing a compound having the formula VIA is provided where R^1 , R^3 , Z^1 , Z^2 , Z^3 , R^4 , R^5 , and R^6 have any of the values described above with respect to compounds of formula IC and embodiments of compounds of formula IC, and Y^1 is selected from the group consisting of NO_2 , a protected amine group, a halogen such as Cl, F, Br, or I, an $-NCS$, and an N_3 group. In some embodiments of the method, R^1 is a substituted or unsubstituted arylalkyl group. In some embodiments of the method, R^3 is H. In some embodiments, Y^1 is selected from a F, NO_2 , or an N_3 group. In some embodiments of the method, R^1 is a substituted or unsubstituted arylalkyl group, R^3 is H, and Y^1 is selected from a F, NO_2 , or an N_3 group. In some such embodiments, Z^1 , Z^2 ,


and Z^3 are each carbon atoms and R^4 , R^5 , and R^6 are each H. Compounds of formula VIA may be readily converted into compounds of formula IC as shown in Scheme 1a and the procedures set forth herein. Typically a compound of formula VIA where Y^1 is a $-NCS$ is reacted with a first amine and then is reacted with a second amine as described in Procedure 1a. Any of the amines defined by the $R^{1'}$, $R^{2'}$, and $R^{3'}$ of compounds of formula IC may be used. The versatility of this procedure allows a wide range of compounds of formula IC to be prepared where W is a guanidine group prepared from compounds of formula VIA where Y^1 is a $-NCS$ group. Another procedure that may be used to prepare compounds of formula IC from compounds of formula VIA where Y^1 is an N_3 group is shown in Scheme 1b where nitrogen compounds with any of the $R^{1'}$, $R^{2'}$, and $R^{3'}$ groups of compounds of formula IC may be used.



VIA


[0195] A method for preparing a compound of formula VIA typically includes cyclizing a compound of formula VIB by reacting it with an electrophilic carbon bearing an R^3 group such as a trialkyl orthoformate such as triethyl orthoformate, trimethyl orthoformate or the like where the R^3 is a H; Gold's Reagent; a substituted or unsubstituted alkanoyl halide such as acetyl chloride where R^3 is a methyl group; a substituted or unsubstituted alkanolic acid in the presence of an acid halide producing agent such as thionyl chloride, $POCl_3$, various phosphorous halides, and the like (e.g. an alkanolic acid of formula R^3CO_2H in combination with $POCl_3$); a benzoyl chloride or an analogous heteroaryl acid chloride compound; or a substituted or

unsubstituted benzoic acid or analogous heteroaryl carboxylic acid compound and an acid halide producing agent. The reaction provides the compound of formula VIA. Compounds of formula VIB have the following formula



VIB

In compounds of formula VIB, the variable may have any of the same values as described above with respect to compounds of formula VIA. The method of forming a compound of formula VIA may include reacting a compound of formula VIC with an amine of formula R^1-NH_2 to produce the compound of formula VIB using standard amide-forming procedures and where R^1 has any of the values of the compounds of formula VIA and VIB and the variables in the compound of VIC have the values of compounds of formula VIA and VIB and Y^2 is a hydroxyl group or is an equivalent thereof. Compounds of formula VIC have the following formula.



VIC

[0196] An alternative method for synthesizing compounds of formula IC is shown for analogous compounds of formula IA in Scheme 2a. Such a method generally includes reacting a compound of formula VIC with an amine



of formula R^1-NH_2 and an electrophilic carbon bearing an R^3 group such as a carboxylic acid of formula R^3-CO_2H . Compounds of formula IC may further be prepared according to a procedure analogous to that shown in Scheme 2b by reacting a compound of formula VIC with an amide of formula $R^1-N(H)-C(=O)-R^3$ in the presence of $POCl_3$ or an analogous compound.

[0197] The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.


EXAMPLES

[0198] The following abbreviations and terms are used throughout the Examples:


Boc:	t-Butyl carbamate protecting group
Celite®:	Diatomaceous earth filter agent
DAST:	(Dimethylamino)sulfur trifluoride
DCM:	Dichloromethane
DIBAL:	Diisobutylaluminum hydride
DIEA:	<i>N,N</i> -Diisopropylethylamine
DMF:	<i>N,N</i> -Dimethylformamide
DMSO:	Dimethylsulfoxide
EDCI:	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtOAc:	Ethyl acetate
EtOH:	Ethanol
Gold's Reagent:	(Dimethylaminomethyleneaminomethylene)dimethyl ammonium chloride
HOBt:	Hydroxybenzotriazole
HPLC:	High performance liquid chromatography
HCl:	Hydrochloric acid
HBTU:	O-Benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
KOH:	Potassium hydroxide
LC:	Liquid Chromatography
MS:	Mass Spectroscopy
MeOH:	Methanol
mL:	Milliliter
NMP:	1-Methyl-2-pyrrolidinone
NMR:	Nuclear magnetic resonance spectroscopy
TFA:	Trifluoroacetic acid
THF:	Tetrahydrofuran

SYNTHESIS OF *cis*-4-FLUOROCYCLOHEXYLAMINE**Step 1. Synthesis of *trans*-(*t*-butoxy)-N-(4-hydroxycyclohexyl)-carboxamide**



[0199] A suspension of *trans*-4-aminocyclohexanol (1 equivalent) in THF (0.1 M) was treated with (Boc)₂O (1 equivalent). The mixture was stirred at room temperature overnight, dissolved in chloroform, and washed with water to yield a solid that was used without further purification.

Step 2. Synthesis of *cis*-(*t*-butoxy)-N-(4-fluorocyclohexyl)carboxamide

[0200] To a solution of (*t*-butoxy)-N-(4-hydroxycyclohexyl)carboxamide (1 equivalent) in CH₂Cl₂ (1 M) cooled to -78°C was added dropwise a solution of DAST (1 equivalent) in CH₂Cl₂ (0.5 M). The mixture was stirred at -78°C for 4 hours, and then allowed to rise to room temperature. The solution was poured into saturated NaHCO₃ and extracted with chloroform, dried, and evaporated. The resulting crude product was purified on silica gel, eluting with ethyl acetate/hexane 5:95.

Step 3. Synthesis of *cis*-4-fluorocyclohexylamine


[0201] A solution of *cis*-(*t*-butoxy)-N-(4-fluorocyclohexyl)carboxamide (6.51 mmol) in CH₂Cl₂ (20 mL) was treated with TFA (10 mL) at room temperature. The reaction mixture was stirred for 2 hours, the solvent was removed *in vacuo*, and the crude product was dissolved in water and washed with chloroform. The acidic aqueous phase was cooled at 0°C and made basic by the addition of solid KOH. The resulting mixture was extracted with CH₂Cl₂, dried and filtered, yielding the title compound which was used without further purification and as a 0.3 M solution in CH₂Cl₂.

SYNTHESIS OF 4,4-DIFLUOROCYCLOHEXYLAMINE**Step 1. Synthesis of N-(4,4-difluorocyclohexyl)(*t*-butoxy)carboxamide**

[0202] A solution of (*t*-butoxy)-N-(4-oxocyclohexyl)carboxamide (2.5 g, 11.7 mmol) in CH₂Cl₂ (45 mL) was treated with a solution of DAST (2.63 mL, 19.93 mmol) in CH₂Cl₂ (6 mL) at room temperature. EtOH (141 µL, 2.3 mmol) was added, and the mixture was stirred at room temperature overnight. The solution was poured into saturated NaHCO₃ and extracted with chloroform, dried, and evaporated to yield a 1:1 mixture of the title compound and (*t*-butoxy)-N-(4-fluorocyclohex-3-enyl)carboxamide. This mixture was


dissolved in CH_2Cl_2 (40 mL) and MeOH (14 mL) and cooled to -78°C . Ozone was bubbled into the solution for 50 minutes until it turned green and Me_2S was added (3 equivalents). The reaction mixture was allowed to warm to room temperature, chloroform was added and the organic phase was washed with water, dried, and evaporated to yield the title compound which was used without further purification.

Step 2. Synthesis of 4,4-difluorocyclohexylamine




[0203] A solution of N-(4,4-difluorocyclohexyl)(t-butoxy)carboxamide (6.51 mmol) in CH_2Cl_2 (20 mL) was treated with TFA (10 mL) at room temperature. The reaction mixture was stirred for 2 hours, the solvent was removed *in vacuo*, and the crude product was dissolved in water and washed with chloroform. The acidic aqueous phase was cooled at 0°C and made basic by the addition of solid KOH. The resulting mixture was extracted with CH_2Cl_2 , dried, and filtered yielding the title compound which was used without further purification as a 0.3 M solution in CH_2Cl_2 .

Procedure 1 Synthesis of 6-fluoro analog of 7-azido-quinazoline-4-one (1)




1

Step 1: Synthesis of (2-amino-4,5-difluorophenyl)-N-[2-(4-fluorophenyl)ethyl]-carboxamide



[0204] To a stirred solution of 4,5-difluoro anthranilic acid (2.0 g, 11.6 mmol) in anhydrous THF (30 mL) was added hydroxybenzotriazole hydrate (HOBT) (1.56 g, 11.6 mmol), diisopropylethyl amine (2.01 mL, 11.6 mmol), and 4-fluorophenylethyl amine (1.52 mL, 11.6 mmol). After all of the HOBT had completely dissolved, EDCI (2.21 g, 11.6 mmol) was added and the resulting orange solution was stirred at room temperature for 16 hours. The solvent was removed, and the residue was chromatographed on silica eluting with 15% EtOAc in hexanes giving the desired benzamide (**2**) as white crystals (3.07 g, 10.4 mmol, 90%).


Step 2: Synthesis of 6,7-difluoro-3-[2-(4-fluorophenyl)ethyl]-3-hydroquinazolin-4-one



[0205] The starting benzamide (**2**) was dissolved in trimethyl orthoformate (20 mL) and heated at 120°C under a stream of nitrogen for 3 hours. The solution was cooled, and the solvent was removed by rotary evaporation. The residue was triturated with hexanes, and the solids collected by filtration, washed with hexanes, and dried on the pump. The formamide intermediate was isolated as a white solid and confirmed by NMR.

This intermediate was suspended in POCl_3 (10 mL) and heated to 140°C for 3 minutes. The reaction was cooled, poured over crushed ice, made slightly alkaline with saturated sodium bicarbonate solution, and extracted with EtOAc. The organic layer was collected and dried over magnesium sulfate. Product (3) was isolated as a white solid (1.94 g, 6.38 mmol, 75% for 2 steps).

Step 3: Synthesis of 7-(azadiazomvinyl)-6-fluoro-3-[2-(4-fluorophenyl)ethyl]-3-hydroquinazolin-4-one



[0206] Difluoroquinazoline (3) (1.46 g, 4.6 mmol) was dissolved in DMSO (10 mL), and sodium azide (3 g, 46.0 mmol) was added. The resulting mixture was heated to 70°C with stirring for 4 hours. The reaction was monitored by NMR. The reaction was cooled and diluted with water, and the resulting precipitate collected by filtration and washed with water. The solid was dissolved in methylene chloride and dried (MgSO_4) in order to remove trace water. Product (1) was isolated as an off-white solid (1.43 g, 4.37 mmol, 95%).

[0207] Following the formation of compound 1, final guanidino quinazilones were formed following the synthetic method described below (Procedure 1A):


Procedure 1A

[0208] To a solution of (1) (1 equivalent) in THF was added trimethylphosphine (1.5 equivalents), and the mixture was stirred at room temperature for 10 minutes. To the iminophosphorane solution was added (1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl isocyanate (1.6


equivalents). The solution was heated at 70°C overnight. To half of the carboimide solution was added a THF solution of (6S,2R)-2,6-dimethylpiperazine (2 equivalents). After being heated at 70°C for 2 hours, the residue was subjected to HPLC purification to give the guanidine product as its TFA salt.

[0209] The 2-fluoro-4-methoxy, 2,4-difluoro and 2,4-dichloro analogs were synthesized via the same pathway described above. Compounds of the group synthesized via the pathway described above include Examples 42, 44, and 45.

EXAMPLE 1. Synthesis of (3R,5S)-N-(3-{2-[2-fluoro-4-(methoxy)-phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide




Step 1. Synthesis of (c): 2-amino-N[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-4-nitro-benzamide



[0210] 2-Fluoro-4-methoxyphenylethylamine ((a): 1 equivalent), 4-nitroanthranilic acid ((b): 1 equivalent), HBTU (1.5 equivalents), and dry THF (0.5 M in (a)) were added to a dry round bottom flask. The mixture was allowed to stir for 10 hours at room temperature. The reaction was then dry loaded onto silica gel and purified via flash chromatography using


hexanes/ethyl acetate. The pure fractions were combined and concentrated *in vacuo* to yield the product ((c): 2-amino-N-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-4-nitro-benzamide) as a pure solid.

Step 2. Synthesis of (d): 3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-7-nitro-3H-quinazolin-4-one




[0211] The pure product ((c): 1 equivalent) of Step 1, Gold's reagent, and dioxane (0.5 M in (c)) were added to a dry round bottom flask, fitted with a condenser, and heated to reflux for 16 hours. Once complete product conversion was verified by LC/MS, acetic acid (1 equivalent) and sodium acetate (1 equivalent) were added to the reaction. The subsequent mixture was heated to reflux for 3 hours. Then, the reaction was concentrated *in vacuo*, taken up in ethyl acetate, and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated. The crude product mixture was purified via flash chromatography using a mixture of $\text{CH}_2\text{Cl}_2/\text{MeOH}$. The pure fractions were combined and concentrated *in vacuo* to yield the pure product ((d): 3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-7-nitro-3H-quinazolin-4-one) as a pure solid.

Step 3. Synthesis of (e): 7-amino-3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-3H-quinazolin-4-one




[0212] To a solution of (d), prepared as described in Step 2, in MeOH (0.25 M in (d)) was added 10% Pd/C (0.1 equivalents). The mixture was sealed with a septum and degassed with nitrogen for 10 minutes. Hydrogen was then bubbled through the solution for 20 minutes. Once reaction completion was verified by LC/MS, the reaction was degassed with nitrogen for 10 minutes. The mixture was filtered through Celite® and concentrated *in vacuo* to yield the product ((e): 7-amino-3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-3H-quinazolin-4-one). The product was used in the next reaction without further purification.

Step 4. Synthesis of (f): 3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-7-isothiocyanato-3H-quinazolin-4-one




[0213] To a mixture of (e), prepared as described in Step 3, (1 equivalent) and NaHCO₃ (3 equivalents) in acetone (0.1 M in (e)) was added thiophosgene (3 equivalents) dropwise. The resulting slurry was stirred at room temperature for three hours. Once reaction completion was verified by LC/MS, the reaction was concentrated *in vacuo* to remove solvent and excess thiophosgene. The mixture was then taken up in ethyl acetate and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated *in vacuo* to yield the product ((f): 3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-7-isothiocyanato-3H-quinazolin-4-one). The crude product was used in the next reaction without further purification.

Step 5. Synthesis of (g): 1-{3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-4-oxo-3,4-dihydro-quinazolin-7-yl}-3-(2,6,6-trimethyl-bicyclo[3.1.1]hept-3-yl)-thiourea



[0214] To a solution of (f), prepared as described in Step 4, (1 equivalent) in THF (0.5 M in (f)) was added (1S,2S,3S,5R)-(+)-isopinocampheylamine (1.5 equivalents). The reaction was stirred at room temperature for 10 hours. The crude product mixture was then concentrated *in vacuo*, dissolved in methylene chloride, and purified via flash chromatography using hexanes/ethyl acetate. The pure fractions were combined and concentrated *in vacuo* to yield the pure product ((g): 1-{3-[2-(2-fluoro-4-methoxy-phenyl)-ethyl]-4-oxo-3,4-dihydro-quinazolin-7-yl}-3-(2,6,6-trimethyl-bicyclo[3.1.1]hept-3-yl)-thiourea).

Step 6. Synthesis of (h): (3R,5S)-N-(3-[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide



[0215] To a solution of (g), prepared as described in Step 5, (1 equivalent) in dry THF (0.1 M in (g)) in a dry round bottom flask was added 1-[3-(dimethylamino)-propyl]-3-ethylcarbodiimide hydrochloride (2 equivalents). The reaction was fitted with a condenser and heated to 80°C for 1 hour. The resulting solution was allowed to cool to room temperature for 20 minutes. A solution of *cis*-2,6-dimethylpiperazine (2 equivalents; 0.5 M in CH₂Cl₂) was then added to the reaction, and the resulting mixture was stirred at room temperature for 10 minutes. The mixture was then diluted with ethyl acetate and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined and concentrated *in vacuo*. The crude mixture was dissolved in DMSO and purified via preparative HPLC using water (0.1% TFA) /acetonitrile (0.1% TFA). The pure fractions were combined and concentrated *in vacuo* to remove the majority of acetonitrile. Sodium carbonate (15 equivalents) was then added to the resulting aqueous solution and the slurry was allowed to sit at room temperature for 1 hour with occasional swirling. The basic aqueous solution was then extracted with 3 separate portions of ethyl acetate. The organic layers were combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated *in vacuo* to yield product (h) as a free base. The resulting solid was then dissolved in an aqueous HCl solution (1 M; 15 equivalents) and concentrated *in vacuo*. The resulting mixture was dissolved in a 1:1 water/acetonitrile mixture and lyophilized to yield the pure Bis-HCl salt product ((h): (3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide.

Synthesis of Compounds of Structure IIIA, IIIB, IIIF, and IIIG

[0216] Compounds of formula IIIA, IIIB, IIIF, and IIIG are prepared using the methodology described above using an appropriately substituted pyridine, pyrazine, or pyrimidine benzoic acid, respectively, in place of the 4-nitroanthranilic acid (b) in Step 1. Steps 2-6 may then be carried out to give

the final product. One skilled in the art will also recognize that the pyridine may be further substituted to produce variously substituted compounds where R^4 , R^5 , and/or R^6 are any of the groups herein described such as, but not limited to, fluoro, chloro, alkyl, and alkaryl.


Synthesis of Compounds of Structure IIIC, IIID, and IIIE

[0217] Compounds of formula IIIC, IIID, and IIIE are prepared using the methodology described above using an appropriately fluorine-substituted 4-nitroanthranilic acid in place of 4-nitroanthranilic acid (**b**) in Step 1. Steps 2-6 may then be carried out to give the final product. One skilled in the art will recognize that a fluorine-substituted 4-nitroanthranilic acid may be used which includes further substituents to produce variously substituted compounds where R^4 , R^5 , and/or R^6 are any of the groups herein described such as, but not limited to, fluoro, chloro, alkyl, and alkaryl.


Synthesis of Compounds of Structure IC

[0218] Compounds of formula IC are prepared using the methodology described above using an appropriately substituted 5-nitroanthranilic acid in place of 4-nitroanthranilic acid (**b**) in Step 1. Steps 2-6 may then be carried out to give the final product. One skilled in the art will recognize that a fluorine-substituted 5-nitroanthranilic acid may be used which includes further substituents to produce variously substituted compounds where R^4 , R^5 , and/or R^6 are any of the groups herein described such as, but not limited to, fluoro, chloro, alkyl, and alkaryl.

EXAMPLE 2. Synthesis of 7-[[1-((5S,3R)-3,5-dimethylpiperaziny)-2-((2S,3S,1R,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)(1Z)-2-azavinyl]amino)-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione




Step 1. Synthesis of (c): 2-amino-N-[2-(2,4-dichlorophenyl)-ethyl]-4-nitro-benzamide




[0219] 2,4-Dichlorophenylethylamine ((a):1 equivalent), 4-nitroanthranilic acid ((b): 1 equivalent), HBTU (1.5 equivalent), and dry THF (0.5 M in (a)) were added to a dry round bottom flask. The mixture was allowed to stir for 10 hours at room temperature. The reaction was then dry loaded onto silica gel and purified via flash chromatography using hexanes/ethyl acetate. The pure fractions were combined and concentrated *in vacuo* to yield the product ((c): 2-amino-N-[2-(2,4-dichloro-phenyl)ethyl]-4-nitrobenzamide) as a pure solid.

Step 2. Synthesis of (d): 3-[2-(2,4-dichlorophenyl)ethyl]-7-nitro-1,3-dihydroquinazoline-2,4-dione




[0220] To a 0.3M solution of (c), prepared as described in Step 1, (2.5 g, 7.5 mmol (c)) in dioxane was added 40 mL of a 20% phosgene solution in toluene, followed by 15 mL triethylamine. After stirring for 1 hour at room temperature, solvent was removed by rotary evaporation followed by high vacuum. The residue was dissolved in ethyl acetate and washed three times with water. After drying with sodium sulfate and rotary evaporation, an orange-brown solid ((d): 3-[2-(2,4-dichlorophenyl)ethyl]-7-nitro-1,3-dihydroquinazoline-2,4-dione) was obtained in over 90% yield.

Step 3. Synthesis of (e): 7-amino-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione




[0221] To a solution of (d), prepared as described in Step 2, in MeOH (0.25 M in (d)) was added 10% Pd/C (0.1 equivalents). The mixture was sealed with a septum and degassed with nitrogen for 10 minutes. Hydrogen was then bubbled through the solution for 20 minutes. Once reaction completion was verified by LC/MS, the reaction was degassed with nitrogen for 10 minutes. The mixture was filtered through Celite® and concentrated *in vacuo* to yield the product ((e)) 7-amino-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione). The product was used in the next reaction without further purification.

Step 4. Synthesis of (f): 3-[2-(2,4-dichlorophenyl)ethyl]-2,4-dioxo-1,3-dihydroquinazolin-7-isothiocyanate




[0222] To a mixture of (e), prepared as described in Step 3, (1 equivalent) and NaHCO_3 (3 equivalents) in acetone (0.1 M in (e)) was added thiophosgene (3 equivalents) dropwise. The resulting slurry was stirred at room temperature for three hours. Once reaction completion was verified by LC/MS, the reaction was concentrated *in vacuo* to remove solvent and excess thiophosgene. The mixture was then taken up in ethyl acetate and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated *in vacuo* to yield the product ((f): 3-[2-(2,4-dichlorophenyl)ethyl]-2,4-dioxo-1,3-dihydroquinazolin-7-isothiocyanate). The crude product was used in the next reaction without further purification.

Step 5. Synthesis of (g): 7-([[(2S,3S,1R,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]amino]thioxomethyl)amino)-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione



[0223] To a solution of (f), prepared as described in Step 4, (1 equivalent) in THF (0.5 M in (f)) was added (1S,2S,3S,5R)-(+)-isopinocampheylamine (1.5 equivalents). The reaction was stirred at room temperature for 10 hours. The crude product mixture was then concentrated *in vacuo*, dissolved in methylene chloride, and purified via flash chromatography using hexanes/ethyl acetate. The pure fractions were combined and concentrated *in vacuo* to yield the pure product ((g 7-([[(2S,3S,1R,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]amino]thioxomethyl)amino)-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione).

Step 6. Synthesis of (h): 7-[[1-((5S,3R)-3,5-dimethylpiperazinyl)-2-((2S,3S,1R,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)(1Z)-2-azavinyl]amino]-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione



[0224] To a solution of (g), prepared as described in Step 5, (1 equivalent) in dry THF (0.1 M in (g)) in a dry round bottom flask was added 1-[3-(dimethylamino)-propyl]-3-ethylcarbodiimide hydrochloride (2 equivalents). The reaction flask was fitted with a water-cooled condenser and heated to 80°C for 1 hour under a nitrogen atmosphere. The resulting solution was cooled to 0°C for 20 minutes. A solution of *cis*-2,6-dimethylpiperazine (2 equivalents; 0.5 M in CH₂Cl₂) was then added to the reaction, and the resulting mixture was stirred at 0°C for 10 minutes. The mixture was then diluted with ethyl acetate and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined and concentrated *in vacuo*. The crude mixture was dissolved in DMSO/acetonitrile and purified via preparative HPLC using water (0.1% TFA) /acetonitrile (0.1% TFA). The pure fractions were combined and concentrated *in vacuo* to remove the majority of


acetonitrile. Sodium hydroxide (10 equivalents) was then added to the resulting aqueous solution and the slurry was allowed to sit at room temperature for 1 hour with occasional swirling. The basic aqueous solution was then extracted with 3 separate portions of ethyl acetate. The organic layers were combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated *in vacuo* to yield product (h) as a free base. The resulting solid was then dissolved in an aqueous HCl solution (1 M; 15 equivalents) and concentrated *in vacuo*. The resulting mixture was dissolved in a 1:1 water/acetonitrile mixture and lyophilized to yield the pure Bis-HCl salt product ((h): 7-[[1-((5S,3R)-3,5-dimethylpiperazinyl)-2-((2S,3S,1R,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)(1Z)-2-azavinyl]amino]-3-[2-(2,4-dichlorophenyl)ethyl]-1,3-dihydroquinazoline-2,4-dione).

Synthesis of Compounds of Structure IVA and IVB


[0225] Compounds of formula IVA and IVB are prepared using the methodology described above using an appropriately substituted pyridine in place of the 4-nitroanthranilic acid (b) in Step 1b. Procedure 1A may then be carried out to give the final product. One skilled in the art will also recognize that the pyridine may be further substituted to produce variously substituted compounds where R⁴, R⁵, and/or R⁶ are any of the groups herein described such as, but not limited to, fluoro, chloro, alkyl, and alkaryl.

[0226] Compounds are prepared using the methodology described above in Procedure A. Procedure 1A may then be carried out to give the final product. One skilled in the art will recognize that a fluorine-substituted 4-nitroanthranilic acid may be used which includes further substituents to produce variously substituted compounds where R⁴, R⁵, and/or R⁶ are any of the groups herein described such as, but not limited to, fluoro, chloro, alkyl, and alkaryl.

EXAMPLE 3. Synthesis of 7-([1-((3S)-3-methylpiperazinyl)(1Z)-2-aza-2-(4,4-difluorocyclohexyl)vinyl]amino)-3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-3-hydroquinazolin-4-one




Step 1. Synthesis of (b): 7-([[(4,4-difluorocyclohexyl)amino]-thioxomethyl]amino)-3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-3-hydroquinazolin-4-one



[0227] To a solution of (a), prepared as (f) described in Step 4 of Example 1, (1 equivalent) in THF (0.5 M in (a)) was added 4,4-difluorocyclohexylamine prepared as described above (1.5 equivalents). The reaction was stirred at room temperature for 10 hours. The crude product mixture was then concentrated *in vacuo*, dissolved in methylene chloride, and purified via flash chromatography using hexanes/ethyl acetate. The pure fractions were combined and concentrated *in vacuo* to yield the pure product

((b): 7-([[(4,4-difluorocyclohexyl)amino]-thioxomethyl]amino)-3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-3-hydroquinazolin-4-one).


Step 2. Synthesis of (c): 7-[[1-((3S)-3-methylpiperazinyl)(1Z)-2-aza-2-(4,4-difluorocyclohexyl)vinyl]amino]-3-[2-(2-fluoro-4-methoxyphenyl)-ethyl]-3-hydroquinazolin-4-one




[0228] To a solution of (b), prepared as described in Step 1, (1 equivalent) in dry THF (0.1 M in (b)) in a dry round bottom flask was added 1-[3-(dimethylamino)-propyl]-3-ethylcarbodiimide hydrochloride (2 equivalents). The reaction was fitted with a condenser and heated to 80°C for 1 hour. The resulting solution was allowed to cool to room temperature for 20 minutes. A solution of (S)-2-methylpiperazine (2 equivalents; 0.5 M in CH₂Cl₂) was then added to the reaction, and the resulting mixture was stirred at room temperature for 10 minutes. The mixture was then diluted with ethyl acetate and washed with water. After the organic layer was isolated, the aqueous layer was extracted with two more portions of ethyl acetate. The organic layers were then combined and concentrated *in vacuo*. The crude mixture was dissolved in DMSO and purified via preparative HPLC using water (0.1% TFA)/acetonitrile (0.1% TFA). The pure fractions were combined and concentrated *in vacuo* to remove the majority of acetonitrile. Sodium

carbonate (15 equivalents) was then added to the resulting aqueous solution and the slurry was allowed to sit at room temperature for 1 hour with occasional swirling. The basic aqueous solution was then extracted with 3 separate portions of ethyl acetate. The organic layers were combined, dried over sodium sulfate, filtered through a cotton plug, and concentrated *in vacuo* to yield product (c) as a free base. The resulting solid was then dissolved in an aqueous HCl solution (1 M; 15 equivalents) and concentrated *in vacuo*. The resulting mixture was dissolved in a 1:1 water/acetonitrile mixture and lyophilized to yield the pure Bis-HCl salt product ((c): 7-[[1-((3S)-3-methylpiperazinyl)(1Z)-2-aza-2-(4,4-difluorocyclohexyl)vinyl]amino]-3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-3-hydroquinazolin-4-one).

Method 1 Synthesis of 3-[2-(4-fluorophenyl)ethyl]-7-nitro-2-(4-pyridyl)-3-hydroquinazolin-4-one




[0229] Pyridine 4-carboxylic acid was stirred in POCl₃ at room temperature for about 5 minutes. To the stirred solution was then added 0.9 equivalents of (2-amino-4-nitrophenyl)-N-[2-(4-fluorophenyl)ethyl]carboxamide. The resulting mixture was then stirred for about 15 minutes at room temperature in a microwave tube, which was then heated to 165°C in a microwave for 10 minutes. LC/MS indicated completion of the reaction. The POCl₃ was evaporated, and the residue was dissolved in CH₂Cl₂ and washed with saturated sodium bicarbonate solution. The combined organic layers were dried over MgSO₄ and concentrated *in vacuo* and chromatographed on silica gel, eluting with a gradient of EtOAc in Hexanes. The resulting product, 3-[2-(4-fluorophenyl)ethyl]-7-nitro-2-(4-pyridyl)-3-hydroquinazolin-4-one, was then converted to Example 77 using the procedures described in Scheme 1a.

Method 2 Synthesis of 2-[2-(2-fluoro-4-methoxyphenyl)ethyl]-3-methyl-7-nitro-3-hydroquinazolin-4-one


[0230] 3-(2-Fluoro-4-methoxy-phenyl)-N-methyl-propionamide was synthesized using an EDCI mediated coupling of 3-(2-fluoro-4-methoxy-phenyl)-N-methyl-propionic acid and methylamine (2M solution in THF). The amide was then taken up in POCl_3 in a microwave vessel and the mixture was stirred about 3 minutes. To the stirred solution was added about 1 equivalent of 4-nitroanthranilic acid. The unsealed vial was stirred for 10 minutes until there was a color change from red to yellow. The vial was then sealed and reacted in a microwave unit at 165°C for 600 seconds. Reaction completion was checked with LC/MS. 2-[2-(2-Fluoro-4-methoxyphenyl)ethyl]-3-methyl-7-nitro-3-hydroquinazolin-4-one was then purified by column chromatography, eluting with EtOAc in hexanes. 2-[2-(2-Fluoro-4-methoxyphenyl)ethyl]-3-methyl-7-nitro-3-hydroquinazolin-4-one was then converted to Example 90 using the procedures described above through the corresponding thiourea (Scheme 1a).

Method 3 Synthesis of 3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-2-(4-methylpiperazinyl)-7-nitro-3-hydroquinazolin-4-one (B) and 3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-2-[imino(4-methylpiperazinyl)-methyl]-7-nitro-3-hydroquinazolin-4-one (C)



[0231] The synthesis of nitrile **A** was first conducted as described in J. Heterocyclic Chem., 35, 659 (1998)). Nitrile **A** was heated in an excess of N-methylpiperazine to 110°C in a microwave for 600 seconds and analyzed by LC/MS to provide **B** and **C**. Products **B** and **C** were separated by column chromatography on silica gel eluting with 10% MeOH in CH₂Cl₂. Compound **B** was the first to come off the column. Compounds **B** and **C** were then respectively converted to Examples 99 and 71 using the procedures described herein.


Method 4 Synthesis of 3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-7-nitro-2-(1,2,3,4-tetraazol-5-yl)-3-hydroquinazolin-4-one




[0232] Nitrile 1 shown above (0.9 g, 2.4 mmoles) was dissolved in dry DMF (5 mL). Sodium azide (0.8 g, 12.2 mmoles) was added and the mixture was heated at 125°C for 1 hour. The reaction was cooled, diluted with water (25 mL), and filtered. The collected solid was redissolved in THF/EtOAc 1:1 (25 mL), washed with water (25 mL), and dried over MgSO₄. Filtration and solvent removal afforded 650 mg of a brown solid. The ¹H NMR (DMSO-d₆, 300 MHz) was consistent with desired product formation. The product was converted to Example 78 using the procedures described herein.

Method 5 Synthesis of 3-[2-(4-fluorophenyl)ethyl]-2-[(4-methylpiperazinyl)-methyl]-7-nitro-3-hydroquinazolin-4-one (3)


Step 1 Synthesis of 2-chloro-N-[2-(4-fluoro-phenyl)-ethyl]-acetamide (1)



[0233] To a solution of 4-fluorophenylethylamine (1.0 equivalent) in dried THF was added Hunig's base (DIEA)(1 equivalent). The mixture was then stirred for 3 minutes at 0°C. Thereafter, a solution of chloroacetylchloride (1.0 equivalent) in THF was added via a syringe over a period of 7 minutes. The reaction mixture was then stirred at room temperature for 1 hour after which time the reaction mixture was condensed *in vacuo*, quenched with water, extracted with ethyl acetate (3X) and dried over Na₂SO₄. After concentration *in vacuo*, compound 1 shown above was obtained, which was carried on further without further purification. LC/MS= M+H 216.1 at 2.18 minutes.

Step 2 Synthesis of 2-chloromethyl-3-[2-(4-fluorophenyl)-ethyl]-7-nitro-3H-quinazolin-4-one (2)


[0234] Compound 1 (1.2 equivalents) was dissolved in neat POCl₃ and allowed to stir under N₂ for 5 minutes. Solid 4-nitroanthranilic acid (1.0 equivalent) was then added, and the mixture was allowed to stir at room temperature for 10 minutes until the color changed to yellow from red. Thereafter, the reaction mixture was refluxed at 100°C for 2 hours, followed by removal of POCl₃ *in vacuo* (addition of triethylamine to the rotovap condenser). The crude product so obtained was neutralized with a saturated solution of NaHCO₃ extracted with ethyl acetate (3 times), dried over Na₂SO₄, and condensed *in vacuo*. Purification of the crude product was carried out with column chromatography in several batches using a gradient of EtOAc in hexanes. LC/MS = M+H 3.62 at 3.5 minutes.

Step 3 Synthesis of 3-[2-(4-fluorophenyl)ethyl]-2-[(4-methylpiperazinyl)-methyl]-7-nitro-3H-quinazolin-4-one (3)

[0235] A solution of 2 (1 equivalent) and 4-methylpiperazine (3 equivalents) in 2 mL NMP were heated at 80°C. After stirring for 18 hours, the dark brown solution was diluted with ethyl acetate and washed twice with water. The organic phase was then dried with sodium sulfate, filtered and concentrated *in vacuo*, and taken on to the next step without further


purification. Compound 3 was then converted to Example 69 using the procedures described herein. This procedure yielded a dark oil, and small amounts of NMP may remain in the product. Formation of some analogous compounds required the addition of three equivalents of diisopropyl ethyl amine. Similar chemistry was used to prepare Examples 67, 70, 72, 74, 75, 79, and 81 as identified in the following tables.

Step 3a Synthesis of 2-[(2,4-difluorophenoxy)methyl]-3-methyl-7-nitro-3-hydroquinazolin-4-one



[0236] 2,4-Difluorophenol (2.5 equivalents) was added to 2-(chloromethyl)-3-methyl-7-nitro-3-hydroquinazolin-4-one (2a) in acetone and refluxed for 8 hours. The solution was then cooled to room temperature, washed with saturated sodium bicarbonate, dried and filtered over sodium sulfate and concentrated *in vacuo* to afford 2-[(2,4-difluorophenoxy)methyl]-3-methyl-7-nitro-3-hydroquinazolin-4-one in quantitative yields. Compound 3a was then converted to Example 88 using the procedures described herein. Similar chemistry was used to prepare Examples 68, 89, 92, 93, 94, 95, 96, 97, 98, and 100 as identified in the following tables.


METHOD 6 Synthesis of 3-[2-(2-fluoro-4-methoxyphenyl)ethyl]-7-nitrobenzo[d]1,2,3-triazin-4-one



[0237] A mixture of benzamide (1) (3.42 mmol), water (40 mL), and concentrated HCl (12 mL) was cooled in an ice bath, and a solution of NaNO₂ (3.6 mmol) in water (5 mL) was added drop wise. The mixture was stirred for 1 hour, and 20 mL 10 N NaOH was added. The stirring was continued for another hour, and the reaction was neutralized with AcOH, extracted with methylene chloride, and dried over MgSO₄. The crude product was chromatographed on silica (30% EtOAc/hexanes) yielding the desired product as a yellow solid. The purified compound was then converted to Example 102 using the procedures described herein.


Method 7 Synthesis of 6-amino-2-[2-(2-fluoro-4-methoxyphenyl)ethyl]-2-hydroisoquinolin-1-one

Step 1



[0238] The diacid **A** (1 equivalent) was added to a flask equipped with a reflux condenser and dean stark trap and charged with dry toluene. The mixture was heated to reflux and then 2-(2-fluoro-4-methoxy-phenyl)-ethylamine **B** (1 equivalent) was added. The reaction was kept at reflux overnight, and then the toluene was removed by rotary evaporation. Purification by flash chromatography using ethyl acetate/hexanes provided the product **C** in 30% yield.

Step 2




[0239] The imide (C) was dissolved in CH₂Cl₂ and cooled to -78°C. 3 equivalents of DIBAL (1M in CH₂Cl₂) were added, and the reaction was stirred at -78°C for 1 hour when LC/MS indicated completion of the reaction. The solution was then diluted with ether and 10 equivalents of NaF and 4 equivalents water were added. The reaction was then stirred for an hour. The reaction was then filtered through Celite® to yield the crude pyridone amine (D). Compound D was then converted to Example 103 using the procedures described herein. Similar chemistry was used to prepare Example 104 as identified in the final table.

[0240] As noted below, the compounds in the following tables were prepared using the methodology described herein from commercially available starting materials which are readily recognizable by those skilled in the art or by using known synthetic methods. For example, Example 11 was prepared using the methodology described in Scheme 1a and the appropriate amino indanol. Examples 14 and 18 which include hydroxymethyl-substituted arylalkyl groups were also prepared using the general methodology of Scheme 1a with the appropriate amino alcohol. N-cyano substituted piperazine compound Example 36 was prepared by: first, mono-Boc protecting 2,6-*trans* dimethylpiperazine; second, treating the mono-Boc protected compound with cyanogen bromide (2.5 equivalents) and Hunig's base (1.1 equivalent); third, purifying the resulting nitrile piperazine compound on silica gel; fourth, deprotecting the purified compound; and fifth, reacting the resulting purified nitrile *trans* dimethyl piperazine compound using the

methods described herein to produce Example 36. Compounds such as Examples 73 and 76 were prepared using the procedure of Method 2 with the appropriated amides of methacrylic acid and acetic acid.







[0241] Compounds of formula IB where R^2 is an alkyl group such as Example 59 where R^2 is a methyl group may be prepared by alkylating a dione where R^2 is H prepared as described herein. For example, Example 59 was prepared using the methylation procedure shown below (reaction of dione with methyl iodide and potassium carbonate (1:2:2) in DMF at 60°C to produce the nitro compound which was then converted to Example 59 using the standard procedures described herein.









[0242] The compounds in the following tables were prepared using the methodology described in Examples 1-3 and the above Methods and Procedures. The starting materials used in the syntheses are recognizable to one of skill in the art and are commercially available or may be prepared using known methods. The synthesis of various guanidine compounds is known in the art. Such synthesis information may be found in the following references each of which is incorporated herein in its entirety: PCT publication WO 02/18327; United States Patent Application Serial No. 09/945,384; United States Provisional Patent Application Serial No. 60/230,565; United States Provisional Patent Application Serial No. 60/245,579; United States Provisional Application Serial No. 60/282,847; United States Provisional Application Serial No. 60/353,183; and United States Provisional Application Serial No. 60/353,188.







Table of Examples 4-66






No.	Structure	Name	MH+
4		(3S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	575.7
5		(3S,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	589.8
6		(3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	589.8
7		(3S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[4-(trifluoromethyl)cyclohexyl]piperazine-1-carboximidamide	589.6
8		N-[2-(dimethylamino)ethyl]-N'-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-N-(phenylmethyl)-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]guanidine	653.9







9		(3S)-N-{3-[2-(4-fluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	545.7
10		(3S)-N-{3-[2-(2,4-difluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	563.7
11		(3S)-N-{3-[(1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	555.7
12		(3S)-N'-(4,4-difluorocyclohexyl)-N-(3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methylpiperazine-1-carboximidamide	557.6
13		(3S)-N'-(4-fluorocyclohexyl)-N-(3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methylpiperazine-1-carboximidamide	539.6
14		(3S)-N-{3-[(1S)-1-(hydroxymethyl)-3-methylbutyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	523.7






15		(3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-(2-methylcyclohexyl)piperazine-1-carboximidamide	549.7
16		(3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-(4-methylcyclohexyl)piperazine-1-carboximidamide	549.7
17		(3S,5S)-N-{3-[2-(2,4-difluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	577.7
18		(3R,5S)-N-{3-[(1S)-2-[2-fluoro-4-(methyloxy)phenyl]-1-(hydroxymethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	619.8
19		(3S,5S)-N-{3-[(1S)-2-[2-fluoro-4-(methyloxy)phenyl]-1-(hydroxymethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	619.8
20		(3R,5S)-N-{3-[2-(4-fluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	559.7






21		(3R)-3-(dimethylamino)-N-(3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]pyrrolidine-1-carboximidamide	589.8
22		(3S)-3-(dimethylamino)-N-(3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]pyrrolidine-1-carboximidamide	589.8
23		(3S)-N-(3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-4-hydroxy-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	591.7
24		(3S,5S)-N-(3-{2-(4-fluorophenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	559.7
25		(3R,5S)-N-(3-{2-(2,4-difluorophenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	577.7
26		(3R,5S)-N-(3-{2-(2-fluoro-4-methylphenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	573.8







27		(3S,5S)-N-{3-[2-(2-fluoro-4-methylphenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	573.8
28		(3R,5S)-N-{3-[(1S)-2-(4-fluorophenyl)-1-(hydroxymethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	589.8
29		(3S)-N-{3-[2-(4-fluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-4-hydroxy-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	561.7
30		(3R,5S)-N-{3-[2-(4-chlorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	576.2
31		(3R,5S)-N-{3-[2-(2,4-dichlorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	610.6
32		(3R,5S)-N-{3-[(1S)-2-(4-chlorophenyl)-1-(hydroxymethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	606.2






33		(3R,5S)-N-{3-[(1S)-2-hydroxy-1-(phenylmethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	571.8
34		(3R,5S)-N-{3-[2-(4-chloro-2-fluorophenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	594.2
35		(3R,5S)-N-{3-[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-3,4-dihydropyrido[2,3-d]pyrimidin-7-yl]-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	590.8
36		(3S,5S)-4-cyano-N-{3-[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	614.8
37		(3S,5S)-N-{3-[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-N,3,5-trimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	603.8



38		(3S)-N-(3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-N'-[(1S,2S,3S,5R)-2-(hydroxymethyl)-6,6-dimethylbicyclo[3.1.1]hept-3-yl]-3-methylpiperazine-1-carboximidamide	591.7
39		(3S)-N-(3-{2-(4-chlorophenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	582.2
40		(3S)-N-(3-[(1S)-2-(2,4-difluorophenyl)-1-(fluoromethyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	595.7
41		(3S)-N-(6-fluoro-3-{2-(4-fluorophenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	563.7
42		(3R,5S)-N-(6-fluoro-3-{2-(4-fluorophenyl)ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	577.7
43		(3S)-N-(3-[(1R)-2-[2-fluoro-4-(methoxy)phenyl]-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	589.8

44		(3S)-N-(6-fluoro-3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	593.7
45		(3R,5S)-N-(6-fluoro-3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	607.8
46		(3S)-N-{3-[2-(2-fluoro-4-methylphenyl)ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	559.7
47		(3R,5S)-N-(3-{(1R)-2-[2-fluoro-4-(methoxy)phenyl]-1-methylethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	603.8
48		(3S,5S)-N-(3-{(1R)-2-[2-fluoro-4-(methoxy)phenyl]-1-methylethyl}-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	603.8

49		(3S)-N-{3-[(1R)-2-(2-fluoro-4-methylphenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	573.8
50		(3S,5S)-N-{3-[(1R)-2-(2-fluoro-4-methylphenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	587.8
51		(3R,5S)-N-{3-[(1R)-2-(2,4-dichlorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	624.7
52		(3S)-N-{3-[(1R)-2-(2,4-dichlorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	610.6
53		(3S)-N-{3-[(1R)-2-(4-fluorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	559.7



54		(3R,5S)-N-{3-[(1R)-2-(4-fluorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	573.8
55		(3S)-N-{3-[(1R)-2-(4-chlorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	576.2
56		(3R,5S)-N-{3-[(1R)-2-(4-chlorophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	590.2
57		(3R,5S)-N-{3-[(1R)-2-(4-bromophenyl)-1-methylethyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	634.7
58		(3R,5S)-N-{3-[2-(2,4-difluorophenyl)ethyl]-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl]-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	593.7
59		(3S)-N-{3-[2-(4-fluorophenyl)ethyl]-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	575.7







60		(3R,5S)-N-{3-[(1S)-2-(2,4-dichlorophenyl)-1-(hydroxymethyl)ethyl]-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	656.7
61		(3S)-N-(2-hydroxy-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	453.6
62		(3S)-N-{3-[2-(4-[(4-fluorophenyl)carbonyl]piperidin-1-yl)ethyl]-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	672.9
63		(3S)-N-{3-[2-(2,4-dichlorophenyl)ethyl]-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl}-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	612.6
64		(3S)-N-(3-{2-[2-fluoro-4-(methoxy)phenyl]ethyl}-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	591.7








65		(3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	605.8
66		(3S)-N'-(4,4-difluorocyclohexyl)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methylpiperazine-1-carboximidamide	573.6







[0243] The compounds in the following table were prepared using the methodology described in the previous Examples and Methods. The starting materials used in the syntheses are recognizable to one of skill in the art and are commercially available or may be prepared using known methods.

Table of Examples 67-101







No.	Structure	Name	MH+
67		(3R,5S)-N-{3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-2-[(4-methylpiperazin-1-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl}-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	701.9
68		(3S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-2-[(methyloxy)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	619.8



69		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-[(4-methylpiperazin-1-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	657.9
70		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(1H-imidazol-1-ylmethyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	625.8
71		(3S)-N-[3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-2-[imino(4-methylpiperazin-1-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	700.9
72		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-[(4-hydroxypiperidin-1-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	658.9
73		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(1-methylethenyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	585.8
74		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-[(3-hydroxypiperidin-1-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	658.9

75		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-(2H-tetrazol-2-ylmethyl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	627.8
76		(3S)-N-(3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-2-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	589.8
77		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-pyridin-4-yl-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	622.8
78		(3S)-N-[3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-4-oxo-2-(2H-tetrazol-5-yl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	643.8
79		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(morpholin-4-ylmethyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	644.8
80		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(1H-imidazol-1-ylmethyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	625.8
81		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-(1H-tetrazol-1-ylmethyl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	627.8

82		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-phenyl-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	621.8
83		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(2-methylpropyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	601.8
84		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-[2-(4-methoxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	679.9
85		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(4-methylcyclohexyl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	641.9
86		(3S)-N-(2,3-bis[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	727.9
87		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-(2-phenylethyl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	649.9



88		(3S)-N-(2-(((2,4-difluorophenyl)oxy)methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-((1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	580.7
89		(3S)-N-[2-(((2-fluoro-4-(methyloxy)phenyl)oxy)methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-((1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	592.7
90		(3S)-N-(2-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-((1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	590.8
91		(3S)-3-methyl-N-[3-methyl-4-oxo-2-(1H-tetrazol-1-yl)methyl]-3,4-dihydroquinazolin-7-yl]-N'-((1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	520.7
92		(3S)-N-(2-(((4-fluorophenyl)oxy)methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-((1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	562.7
93		(3S)-N-(2-(((4-chlorophenyl)oxy)methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-((1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl)piperazine-1-carboximidamide	579.2







94		(3R,5S)-N-(2-([(4-fluorophenyl)oxy]methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	576.7
95		(3R,5S)-N-(2-([(2,4-difluorophenyl)oxy]methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	594.7
96		(3R,5S)-N-(2-([(2-fluoro-4-(methyloxy)phenyl]oxy]methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	606.8
97		(3S)-N-(2-([(2,4-dichlorophenyl)oxy]methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	613.6
98		(3R,5S)-N-(2-([(2,4-dichlorophenyl)oxy]methyl)-3-methyl-4-oxo-3,4-dihydroquinazolin-7-yl)-3,5-dimethyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	627.6
99		(3S)-N-[3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-2-(4-methylpiperazin-1-yl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	673.9




100		(3S)-N-[2-({[2-fluoro-4-(methyloxy)phenyl]oxy)methyl}-4-oxo-3-(phenylmethyl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	667.8
101		(3S)-3-methyl-N-{2-[(4-methylpiperazin-1-yl)methyl]-4-oxo-3-prop-2-enyl-3,4-dihydroquinazolin-7-yl}-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	575.8

[0244] The compounds in the following table were prepared using the methodology described in the previous Examples and Methods. The starting materials used in the syntheses are recognizable to one of skill in the art and are commercially available or may be prepared using known methods.

Table of Examples 102-112

No.	Structure	Name	MH+
102		(3R,5S)-N-(3-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-4-oxo-3,4-dihydro-1,2,3-benzotriazin-7-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	590.8
103		(3R,5S)-N-(2-{2-[2-fluoro-4-(methyloxy)phenyl]ethyl}-1-oxo-1,2-dihydroisoquinolin-6-yl)-3,5-dimethyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	588.8

104		(3S)-N-[2-[2-(2,4-difluorophenyl)ethyl]-1-oxo-1,2-dihydroisoquinolin-6-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	562.7
105		(3S)-N-[3-[2-[2-fluoro-4-(methoxy)phenyl]ethyl]-4-oxo-2-(1,3-thiazol-2-yl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	658.9
106		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-(4H-1,2,4-triazol-3-yl)-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	612.8
107		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-pyridin-3-yl-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	622.8
108		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-pyridin-2-yl-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	622.8
109		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-4-oxo-2-pyrazin-2-yl-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	623.8

110		(3S)-N-[3-[2-(4-fluorophenyl)ethyl]-2-(5-methylpyrazin-2-yl)-4-oxo-3,4-dihydroquinazolin-7-yl]-3-methyl-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	637.8
111		(3R)-3-(fluoromethyl)-N-(3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	593.7
112		(3R)-N-(3-[2-[2-fluoro-4-(methyloxy)phenyl]ethyl]-4-oxo-3,4-dihydroquinazolin-7-yl)-3-(trifluoromethyl)-N'-[(1S,2S,3S,5R)-2,6,6-trimethylbicyclo[3.1.1]hept-3-yl]piperazine-1-carboximidamide	629.7

[0245] EC₅₀ values of test compounds were determined by treating cells expressing MC4-R with test compound and lysing the cells and measuring intercellular cAMP concentration with an Amersham-Pharmacia RPA-559 cAMP Scintillation Proximity Assay (SPA) kit. The compounds described above were synthesized and tested according to this assay. Each of the named compounds of Examples 1-104 exhibited -log EC₅₀ values above about 3. Examples 105-112 will be found to exhibit -log EC₅₀ values above about 3. For this reason, each of the exemplary compounds are individually preferred and are preferred as a group. Furthermore, the groups corresponding to R¹ through R⁹, R^{1'} through R^{4'}, L, Y, and W for each of the named compounds of Examples 1-112 are also preferred. Nomenclature for these compounds was provided using ACD Name version 5.07 software (November 14, 2001) available from Advanced Chemistry Development, Inc and ChemInnovation NamExpert + Nomenclator™ brand software available from ChemInnovation Software, Inc. Some of the starting materials were

named using standard IUPAC nomenclature. Example compounds 1-112 are illustrative and should not be construed as limiting of the instant invention.

In Vivo Studies of MC4-R Agonists on Energy Intake, Body Weight, Hyperinsulinemia, and Glucose Levels

[0246] In vivo studies are conducted to observe the effect of MCR-4 agonists on energy intake, body weight, hyperinsulinemia, and glucose levels. All studies are conducted with male 9-10 week old ob/ob mice which display early onset of obesity, insulin resistance and diabetes due to leptin deficiency. Mice are acclimated in the facility for 1 week before studies and are caged individually. Vehicle-treated (control) and drug treated mice studies are always run in parallel. In multi-day studies, mice (8-15 per group) are monitored for baseline body weight, fasting levels of glucose, insulin, blood lipids and energy expenditure and then injected twice daily (9 a.m. and 5 p.m.) with 3 mg/kg of a MC4-R agonist of the present invention for 4 weeks. Body weight as well as food and water intake are monitored daily. Animals are fasted overnight for measurements of fasting levels of glucose, insulin, and lipids once a week until the end of the study. Energy expenditure (resting metabolic rate, i.e., O₂ consumption and CO₂ production) are monitored in air tight chambers at the end of the study on fed animals. O₂ consumption and CO₂ production are measured using Oxymax systems (Columbus Instruments). Oral glucose tolerance test (OGTT – a routine test for diabetes and glucose intolerance) is performed on overnight fasted mice at the end of the study. Blood glucose and oral glucose tolerance are measured using a glucose monitor (Onetouch sold by Lifescan). Free fatty acids are measured using an non-esterified free fatty acids enzymatic assay (Waco Chemicals). Serum Insulin levels are measured by immunoassay (Alpco).

Results

[0247] The effect of the compounds of the present invention on food intake is determined by measuring grams/mouse/day throughout a 4 week

study. Food is monitored every morning. Cumulative food intake represents the total amount of grams the mice consume during the study. A significant reduction in food intake is demonstrated in those mice treated IP with the compounds of the present invention.

[0248] The effect of the compounds of the present invention on body weight is determined by measuring grams/mouse throughout a 4 week study. Mice are weighed every morning. A significant body weight reduction is demonstrated in those mice treated IP with the compounds of the present invention.

[0249] The effect of the compounds of the present invention on blood glucose levels is determined by measuring blood glucose levels as represented as mg of glucose/dL of blood. Mice are fasted overnight and glucose levels are measured the following morning. Vehicle treated mice show an increase in blood glucose consistent with the rapid progression of diabetes in this mouse strain whereas, diabetes is slowed down considerably in drug treated mice. A significant reduction in fasting glucose levels is demonstrated in those mice treated IP with the compounds of this invention.

[0250] The effect of the compounds of the present invention on glucose levels during oral glucose tolerance test (OGTT) is determined by measuring blood glucose in overnight fasted mice. Blood glucose is represented as mg of glucose/dL of blood. Glucose levels are measured the following morning. Orally administered glucose quickly elevates blood glucose, similar to a meal, and the response to this exogenous glucose gives a measure of how well the body regulated glucose homeostasis. Vehicle treated mice show an elevated response to glucose consistent with their diabetic state, whereas drug treated mice show a very much improved glucose disposal.

[0251] The effect of the compounds of the present invention on free fatty acid (FFA) levels is determined by measuring mmoles of FFA/L of serum. Mice are fasted overnight and free fatty acid levels are measured the following

morning. Vehicle treated mice show elevated levels of FFA throughout the study consistent with their obese state, whereas the drug treated mice diabetes show a dramatic decrease.


[0252] The effect of the compounds of the present invention on serum insulin levels is determined by measuring serum insulin levels one hour after single IP dosing of 1 and 3 mg/kg in overnight fasted ob/ob mice. Serum insulin levels are represented as ng of insulin/mL of serum. Drug treated mice show a dose dependent decrease relative to vehicle.

[0253] It is understood that the invention is not limited to the embodiments set forth herein for illustration, but embraces all such forms thereof as come within the scope of the following claims.


CLAIMS

What is claimed is:


1. A compound of formula IA, IB, or IC



IA



IB



IC

2

3 wherein

4 Z^1 is selected from the group consisting of CR^4 and N;

5 Z^2 is selected from the group consisting of CR^5 and N;

6 Z^3 is selected from the group consisting of CR^6 and N;

7 R^1 is selected from the group consisting of substituted and
8 unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl,
9 cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;


10 R^2 is selected from the group consisting of H, and substituted
11 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,

12 heterocyclyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, alkylcarbonyl, and
13 arylcarbonyl groups;


14 R^3 is selected from the group consisting of H, and substituted
15 and unsubstituted arylalkyl, heteroarylalkyl, alkoxy, alkylamino, dialkylamino,
16 aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl,
17 alkenyl, alkynyl, and alkyl groups;

18 R^4 , R^5 , and R^6 are independently selected from the group
19 consisting of H, Cl, I, F, Br, OH, NH₂, CN, NO₂, and substituted and
20 unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino,
21 cycloalkyl, heterocyclylamino, heteroarylamino, aminocarbonyl,
22 alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl,
23 arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl
24 groups;

25 W is a group of formula IIA or IIB;



IIA



IIB

26 $R^{1'}$ is selected from the group consisting of H, and substituted
27 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
28 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

30 $R^{2'}$ is selected from the group consisting of H, and substituted
31 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
32 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

33 or R^{1'} and R^{2'}, together with the nitrogen to which they are
34 bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

35 R^{3'} is selected from the group consisting of H, and substituted
36 and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,
37 heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl
38 groups;

39 R^{4'} is selected from the group consisting of H, and substituted
40 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl,
41 cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl
42 groups; and

43 prodrugs thereof, pharmaceutically acceptable salts thereof,
44 stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof,
45 or solvates thereof.

1 2. The compound of claim 1, wherein R³ is H.

1 3. The compound of claim 1, wherein Z¹ is a CR⁴ group, Z²
2 is a CR⁵ group, and Z³ is a CR⁶ group.

1 4. The compound of claim 3, wherein at least one of R⁴, R⁵,
2 or R⁶ is a F.

1 5. The compound of claim 1, wherein at least one of Z¹, Z²,
2 or Z³ is N.

1 6. The compound of claim 1, wherein R^{3'} is selected from
2 the group consisting of substituted and unsubstituted cycloalkyl, polycyclic
3 cycloalkyl, alkenyl, alkyl, and aryl groups.

1 7. The compound of claim 1, wherein R^{1'} is H and R^{2'} is
2 selected from the group consisting of substituted and unsubstituted alkyl,
3 arylalkyl, and heteroarylalkyl groups.

- 1 8. The compound of claim 1, wherein $R^{1'}$ is H and $R^{2'}$ is
2 selected from the group consisting of substituted and unsubstituted
3 dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-
4 methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and
5 thiophene groups.
- 1 9. The compound of claim 1, wherein $R^{1'}$ and $R^{2'}$ may be the
2 same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl
4 groups.
- 1 10. The compound of claim 1, wherein $R^{1'}$ and $R^{2'}$ may be the
2 same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl,
4 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-
5 methoxybenzyl, 2-chlorobenzyl, and thiophene groups.
- 1 11. The compound of claim 1, wherein $R^{1'}$ and $R^{2'}$, together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 heterocyclyl group.
- 1 12. The compound of claim 1, wherein $R^{1'}$ and $R^{2'}$, together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino
4 group.
- 1 13. The compound of claim 1, wherein R^1 is a 2,4-
2 disubstituted phenylethyl group.
- 1 14. The compound of claim 1, wherein R^1 is selected from the
2 group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-
3 methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-
4 methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl,
5 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-

6 methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, indolyethyl, 4-
7 hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-
8 fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-
9 bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-
10 fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-
11 dimethoxyphenylethyl, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, and (phenyl)(hydroxymethyl)ethyl groups.

1 15. The compound of claim 1, wherein R^1 is selected from the
2 group consisting of substituted and unsubstituted
3 (heteroaryl)(hydroxymethyl)ethyl, (aryl)(hydroxymethyl)ethyl groups,
4 (aryl)(alkoxymethyl)ethyl, (aryl)(aryloxymethyl)ethyl,
5 (aryl)(arylalkoxymethyl)ethyl, (aryl)(heteroaryloxymethyl)ethyl,
6 (aryl)(heterocyclyloxymethyl)ethyl, (heteroaryl)(alkoxymethyl)ethyl,
7 (heteroaryl)(aryloxymethyl)ethyl, (heteroaryl)(arylalkoxymethyl)ethyl,
8 (heteroaryl)(heteroaryloxymethyl)ethyl, and
9 (heteroaryl)(heterocyclyloxymethyl)ethyl groups.


1 16. The compound of claim 1, wherein the compound has the
2 formula IB and R^2 is H.

1 17. A composition comprising the compound of claim 1 and a
2 pharmaceutically acceptable carrier.

1 18. A method of treating an MC4-R mediated disease,
2 comprising administering to a subject in need thereof, the compound of claim
3 1.

1 19. The method according to claim 18, wherein the disease is
2 obesity or type II diabetes.

1 20. A compound of formula IA or IC



3 wherein

4 Z^1 is selected from the group consisting of CR^4 and N;

5 Z^2 is selected from the group consisting of CR^5 and N ;


6 Z^3 is selected from the group consisting of CR^6 and N;

R¹ is selected from the group consisting of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;


10 **R³** is selected from the group consisting of substituted and
11 unsubstituted arylalkyl, heteroarylalkyl, alkoxy, aryloxyalkyl, alkylamino,
12 dialkylamino, aryl, heteroaryl, heterocyclyl, cycloalkyl, heterocyclylalkyl,
13 cycloalkylalkyl, alkenyl, alkynyl, alkyl groups, and -C(=NH)-heterocyclyl
14 groups, and groups of formula -LR⁷;

R⁴, R⁵, and R⁶ are independently selected from the group consisting of H, Cl, I, F, Br, OH, NH₂, CN, NO₂, and substituted and unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino, cycloalkyl, heterocyclylamino, heteroaryl amino, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl, arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl groups;

22 W is a group of formula IIA or IIB;



IIA



IIB

23

24 R^{1'} is selected from the group consisting of H, and substituted
25 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
26 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

27 R^{2'} is selected from the group consisting of H, and substituted
28 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
29 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;


30 or R^{1'} and R^{2'}, together with the nitrogen to which they are
31 bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

32 R^{3'} is selected from the group consisting of H, and substituted
33 and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,
34 heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl
35 groups;

36 R^{4'} is selected from the group consisting of H, and substituted
37 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl,
38 cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl
39 groups;

40 L is selected from the group consisting of a covalent bond,
41 -CH₂-, -O-, -S-, and -NH-;

42 R^7 is selected from the group consisting of substituted and
 43 unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from
 44 a group of formula IIC;



IIC

45
 46 wherein Y is selected from the group consisting of CH₂, O, S,
 47 and NR⁹ where R⁹ is selected from the group consisting of H, and substituted
 48 and unsubstituted alkyl groups;

49 R⁸ is selected from the group consisting of H, a halogen,
 50 hydroxyl, carboxylic acid, and substituted and unsubstituted alkyl, amino,
 51 alkylamino, dialkylamino, alkylaminoalkyl, heterocyclyl, alkoxy, carbonyl, and
 52 aminocarbonyl groups;

53 m is an integer selected from 0, 1, or 2; and

54 prodrugs thereof, pharmaceutically acceptable salts thereof,
 55 stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof,
 56 or solvates thereof.

1 21. The compound of claim 20, wherein R¹ is H.

1 22. The compound of claim 20, wherein R¹ is an alkyl group.

1 23. The compound of claim 20, wherein R¹ is an alkyl group
 2 selected from the group consisting of methyl, ethyl, propyl, butyl, and pentyl
 3 groups.

1 24. The compound of claim 20, wherein R¹ is a substituted or
 2 unsubstituted alkenyl group.

- 1 25. The compound of claim 20, wherein Z^1 is a CR^4 group, Z^2
2 is a CR^5 group, and Z^3 is a CR^6 group.
- 1 26. The compound of claim 25, wherein at least one of R^4 ,
2 R^5 , or R^6 is a F.
- 1 27. The compound of claim 20, wherein at least one of Z^1 , Z^2 ,
2 or Z^3 is N.
- 1 28. The compound of claim 20, wherein R^1 is H and R^2 is
2 selected from the group consisting of substituted and unsubstituted alkyl,
3 arylalkyl, and heteroarylalkyl groups.
- 1 29. The compound of claim 20, wherein R^1 is H and R^2 is
2 selected from the group consisting of substituted and unsubstituted
3 dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-
4 methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and
5 thiophene groups.
- 1 30. The compound of claim 20, wherein R^1 and R^2 may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl
4 groups.
- 1 31. The compound of claim 20, wherein R^1 and R^2 may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl,
4 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-
5 methoxybenzyl, 2-chlorobenzyl, and thiophene groups.
- 1 32. The compound of claim 20, wherein R^1 and R^2 , together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 heterocyclyl group.

1 33. The compound of claim 20, wherein R^{1'} and R^{2'}, together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino
4 group.

1 34. The compound of claim 20, wherein R¹ is a 2,4-
2 disubstituted phenylethyl group.

1 35. The compound of claim 20, wherein R¹ is selected from
2 the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-
3 methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-
4 methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl,
5 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-
6 methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl,
7 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-
8 methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl,
9 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-
10 difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl
11 groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and
13 unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
14 (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
15 (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
16 (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted
17 (aryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted
18 (aryl)(heteroaryloxyethyl)ethyl, substituted and unsubstituted
19 (aryl)(heterocyclyloxyethyl)ethyl, substituted and unsubstituted
20 (heteroaryl)(alkoxyethyl)ethyl, substituted and unsubstituted
21 (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted
22 (heteroaryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted
23 (heteroaryl)(heteroaryloxyethyl)ethyl, and substituted and unsubstituted
24 (heteroaryl)(heterocyclyloxyethyl)ethyl groups.

1 36. The compound of claim 20, wherein R^3 is a substituted or
2 unsubstituted alkenyl group.

1 37. The compound of claim 20, wherein R^3 is a 2,4-
2 disubstituted phenylethyl group.


1 38. The compound of claim 20, wherein R^3 is selected from
2 the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-
3 methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-
4 methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl,
5 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-
6 methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl,
7 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-
8 methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl,
9 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-
10 difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl
11 groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and
13 unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
14 (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
15 (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
16 (aryl)(aryloxyethyl)ethyl, substituted and unsubstituted
17 (aryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted
18 (aryl)(heteroaryloxyethyl)ethyl, substituted and unsubstituted
19 (aryl)(heterocyclyloxyethyl)ethyl, substituted and unsubstituted
20 (heteroaryl)(alkoxyethyl)ethyl, substituted and unsubstituted
21 (heteroaryl)(aryloxyethyl)ethyl, substituted and unsubstituted
22 (heteroaryl)(arylalkoxyethyl)ethyl, substituted and unsubstituted
23 (heteroaryl)(heteroaryloxyethyl)ethyl, and substituted and unsubstituted
24 (heteroaryl)(heterocyclyloxyethyl)ethyl groups.

- 1 39. The compound of claim 20, wherein R^3 is a substituted
2 alkyl group selected from substituted or unsubstituted aryloxyalkyl groups or
3 substituted or unsubstituted heteroaryloxyalkyl groups.
- 1 40. The compound of claim 20, wherein R^3 is a substituted or
2 unsubstituted $-CH_2-O$ -aryl group.
- 1 41. The compound of claim 40, wherein the $R^3 -CH_2-O$ -aryl
2 group is substituted with at least one halogen group.
- 1 42. The compound of claim 40, wherein the $R^3 -CH_2-O$ -aryl
2 group is substituted with at least one alkoxy group.
- 1 43. The compound of claim 20, wherein R^3 is a $-CH_2-O$ -aryl
2 group where the aryl group is selected from 2,4-difluorophenyl,
3 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-methoxyphenyl, 2,4-dichlorophenyl,
4 4-chlorophenyl, 2-chlorophenyl, or 2-chloro-4-methoxyphenyl groups.
- 1 44. The compound of claim 20, wherein R^3 is a substituted or
2 unsubstituted heterocyclalkyl group.
- 1 45. The compound of claim 20, wherein R^3 is a substituted or
2 unsubstituted $-CH_2$ -heterocycl group.
- 1 46. The compound of claim 45, wherein the heterocycl
2 group of the $R^3 -CH_2$ -heterocycl group is selected from the group consisting
3 of substituted and unsubstituted 1H-tetrazole, piperazine, piperidine,
4 imidazole, and morpholine groups.
- 1 47. The compound of claim 20, wherein R^3 is selected from a
2 substituted or unsubstituted heterocycl group, a substituted or unsubstituted
3 cycloalkyl group, a substituted or unsubstituted phenyl group, or a substituted
4 or unsubstituted alkoxyalkyl group.
- 1 48. A composition comprising the compound of claim 20 and
2 a pharmaceutically acceptable carrier.

1 49. A method of treating an MC4-R mediated disease,
 2 comprising administering to a subject in need thereof, the compound of claim
 3 20.

1 50. The method according to claim 49, wherein the disease is
 2 obesity or type II diabetes.

1 51. A compound of formula ID




2
 3 wherein

4 R¹ is selected from the group consisting of H, substituted and
 5 unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocyclyl,
 6 cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;


7 R² and R³ are independently selected from the group consisting
 8 of H, substituted and unsubstituted arylalkyl, heteroarylalkyl, alkoxy,
 9 aryloxyalkyl, alkylamino, dialkylamino, aryl, heteroaryl, heterocyclyl,
 10 cycloalkyl, heterocyclylalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkyl groups, and
 11 -C(=NH)-heterocyclyl groups, and groups of formula -LR⁷;

12 R⁴, R⁵, and R⁶ are independently selected from the group
 13 consisting of H, Cl, I, F, Br, OH, NH₂, CN, NO₂, and substituted and
 14 unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino,
 15 cycloalkyl, heterocyclylamino, heteroarylamino, aminocarbonyl,
 16 alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl,
 17 arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl
 18 groups;

19 **W** is a group of formula IIA or IIB;



IIA



IIB

20

21 **R^{1'}** is selected from the group consisting of H, and substituted
 22 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
 23 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

24 **R^{2'}** is selected from the group consisting of H, and substituted
 25 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
 26 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;


27 or **R^{1'}** and **R^{2'}**, together with the nitrogen to which they are
 28 bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

29 **R^{3'}** is selected from the group consisting of H, and substituted
 30 and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,
 31 heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl
 32 groups;

33 **R^{4'}** is selected from the group consisting of H, and substituted
 34 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl,
 35 cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl
 36 groups;

37 **L** is selected from the group consisting of a covalent bond,
 38 -CH₂-, -O-, -S-, and -NH-;

39 R^7 is selected from the group consisting of substituted and
 40 unsubstituted arylaminoalkyl, aryl, and aryloxyalkyl groups or is selected from
 41 a group of formula IIC;



IIC

42
 43 wherein Y is selected from the group consisting of CH₂, O, S,
 44 and NR⁹ where R⁹ is selected from the group consisting of H, and substituted
 45 and unsubstituted alkyl groups;

46 R^8 is selected from the group consisting of H, a halogen,
 47 hydroxyl, carboxylic acid, and substituted or unsubstituted alkyl, amino,
 48 alkylamino, dialkylamino, alkylaminoalkyl, heterocyclyl, alkoxy, carbonyl, and
 49 aminocarbonyl groups;

50 m is an integer selected from 0, 1, or 2; and

51 prodrugs thereof, pharmaceutically acceptable salts thereof,
 52 stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof,
 53 or solvates thereof.

1 52. The compound of claim 51, wherein R^1 is H.

1 53. The compound of claim 51, wherein R^1 is an alkyl group.

1 54. The compound of claim 51, wherein R^2 is H.

1 55. The compound of claim 51, wherein R^2 is a substituted or
 2 unsubstituted alkyl group.

1 56. The compound of claim 51, wherein at least one of R^4 ,
 2 R^5 , or R^6 is a F.

1 57. The compound of claim 51, wherein R^3 is selected from
2 the group consisting of substituted and unsubstituted cycloalkyl, polycyclic
3 cycloalkyl, alkenyl, alkyl, and aryl groups.

1 58. The compound of claim 51, wherein R^1 is H and R^2 is
2 selected from the group consisting of substituted and unsubstituted alkyl,
3 arylalkyl, and heteroarylalkyl groups.

1 59. The compound of claim 51, wherein R^1 is H and R^2 is
2 selected from the group consisting of substituted and unsubstituted
3 dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-
4 methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and
5 thiophene groups.

1 60. The compound of claim 51, wherein R^1 and R^2 may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl
4 groups.

1 61. The compound of claim 51, wherein R^1 and R^2 may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl,
4 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-
5 methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

1 62. The compound of claim 51, wherein R^1 and R^2 , together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 heterocyclyl group.

1 63. The compound of claim 51, wherein R^1 and R^2 , together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino
4 group.

1 64. The compound of claim 51, wherein R¹ is a 2,4-
2 disubstituted phenylethyl group.

1 65. The compound of claim 51, wherein R¹ is selected from
2 the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-
3 methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-
4 methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl,
5 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-
6 methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl,
7 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-
8 methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl,
9 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-
10 difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl
11 groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and
13 unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
14 (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
15 (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
16 (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted
17 (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
18 (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted
19 (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted
20 (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted
21 (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted
22 (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
23 (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted
24 (heteroaryl)(heterocyclyloxymethyl)ethyl groups.

1 66. The compound of claim 51, wherein R¹ is a substituted or
2 unsubstituted alkenyl group.

1 67. The compound of claim 51, wherein R³ is a substituted or
2 unsubstituted alkenyl group.

1 68. The compound of claim 51, wherein R^3 is a 2,4-
2 disubstituted phenylethyl group.

1 69. The compound of claim 51, wherein R^3 is selected from
2 the group consisting of phenylethyl, 2,4-dichlorophenylethyl,
3 4-methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl,
4 4-methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl,
5 cyclohexenylethyl, 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-
6 fluorophenylethyl, 3-methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-
7 hydroxyphenylethyl, 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-
8 fluoro-4-methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-
9 bromophenylethyl, 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-
10 fluorophenylethyl, 2,4-difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-
11 dimethoxyphenylethyl groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and
13 unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
14 (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
15 (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
16 (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted
17 (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
18 (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted
19 (aryl)(heterocyclyloxymethyl)ethyl, substituted and unsubstituted
20 (heteroaryl)(alkoxymethyl)ethyl, substituted and unsubstituted
21 (heteroaryl)(aryloxymethyl)ethyl, substituted and unsubstituted
22 (heteroaryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
23 (heteroaryl)(heteroaryloxymethyl)ethyl, and substituted and unsubstituted
24 (heteroaryl)(heterocyclyloxymethyl)ethyl groups.

1 70. The compound of claim 51, wherein R^3 is a substituted
2 alkyl group selected from substituted or unsubstituted aryloxyalkyl groups or
3 substituted or unsubstituted heteroaryloxyalkyl groups.

1 71. The compound of claim 51, wherein R^3 is a substituted or
2 unsubstituted heterocyclalkyl group.


1 72. The compound of claim 51, wherein R^3 is selected from a
2 substituted or unsubstituted heterocycl group, a substituted or unsubstituted
3 cycloalkyl group, a substituted or unsubstituted phenyl group, or a substituted
4 or unsubstituted alkoxyalkyl group.

1 73. A composition comprising the compound of claim 51 and
2 a pharmaceutically acceptable carrier.

1 74. A method of treating an MC4-R mediated disease,
2 comprising administering to a subject in need thereof, the compound of claim
3 51.

1 75. The method according to claim 74, wherein the disease is
2 obesity or type II diabetes.

1 76. A compound of formula IE




2
3 wherein

4 R^1 is selected from the group consisting of H, substituted and
5 unsubstituted arylalkyl, heteroarylalkyl, aryl, heteroaryl, heterocycl,yl,
6 cycloalkyl, heterocyclalkyl, cycloalkylalkyl, alkenyl, alkynyl, and alkyl groups;


7 R^4 , R^5 , and R^6 are independently selected from the group
8 consisting of H, Cl, I, F, Br, OH, NH_2 , CN, NO_2 , and substituted and

9 unsubstituted alkoxy, alkyl, alkenyl, alkynyl, alkylamino, dialkylamino,
 10 cycloalkyl, heterocyclylamino, heteroaryl amino, aminocarbonyl,
 11 alkylaminocarbonyl, dialkylaminocarbonyl, cycloalkylaminocarbonyl,
 12 arylaminocarbonyl, heterocyclylaminocarbonyl, and heteroarylaminocarbonyl
 13 groups;

14 **W** is a group of formula IIA or IIB;



IIA



IIB

15
 16 **R^{1'}** is selected from the group consisting H, and substituted and
 17 unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, heterocyclyl,
 18 arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

19 **R^{2'}** is selected from the group consisting of H, and substituted
 20 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl,
 21 heterocyclyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl groups;

22 or **R^{1'}** and **R^{2'}**, together with the nitrogen to which they are
 23 bound, form a substituted or unsubstituted heterocyclyl or heteroaryl group;

24 **R^{3'}** is selected from the group consisting of H, and substituted
 25 and unsubstituted aryl, alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl,
 26 heterocyclyl, heterocyclylalkyl, arylalkyl, heteroarylalkyl, and cycloalkylalkyl
 27 groups;

28 **R^{4'}** is selected from the group consisting of H, and substituted
 29 and unsubstituted alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclylalkyl,

30 cycloalkylalkyl, aryl, heteroaryl, heterocyclyl, arylalkyl, and heteroarylalkyl
31 groups; and

32 prodrugs thereof, pharmaceutically acceptable salts thereof,
33 stereoisomers thereof, tautomers thereof, hydrates thereof, hydrides thereof,
34 or solvates thereof.

1 77. The compound of claim 76, wherein R^1 is H.

1 78. The compound of claim 76, wherein R^1 is an alkyl group.

1 79. The compound of claim 76, wherein at least one of R^4 ,
2 R^5 , or R^6 is a F.

1 80. The compound of claim 76, wherein R^3 is selected from
2 the group consisting of substituted and unsubstituted cycloalkyl, polycyclic
3 cycloalkyl, alkenyl, alkyl, and aryl groups.

1 81. The compound of claim 76, wherein $R^{1'}$ is H and $R^{2'}$ is
2 selected from the group consisting of substituted and unsubstituted alkyl,
3 arylalkyl, and heteroarylalkyl groups.

1 82. The compound of claim 76, wherein $R^{1'}$ is H and $R^{2'}$ is
2 selected from the group consisting of substituted and unsubstituted
3 dialkylaminoethyl, 4-ethylbenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-
4 methylbenzyl, benzyl, 4-fluorobenzyl, 3-methoxybenzyl, 2-chlorobenzyl, and
5 thiophene groups.

1 83. The compound of claim 76, wherein $R^{1'}$ and $R^{2'}$ may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted alkyl, arylalkyl, and heteroarylalkyl
4 groups.

1 84. The compound of claim 76, wherein $R^{1'}$ and $R^{2'}$ may be
2 the same or different and are each independently selected from the group
3 consisting of substituted and unsubstituted dialkylaminoethyl, 4-ethylbenzyl,

4 3-chlorobenzyl, 2,4-dichlorobenzyl, 3-methylbenzyl, benzyl, 4-fluorobenzyl, 3-
5 methoxybenzyl, 2-chlorobenzyl, and thiophene groups.

1 85. The compound of claim 76, wherein $R^{1'}$ and $R^{2'}$, together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 heterocyclyl group.

1 86. The compound of claim 76, wherein $R^{1'}$ and $R^{2'}$, together
2 with the nitrogen to which they are bound, form a substituted or unsubstituted
3 piperazino, morpholino, pyrrolidino, piperidino, homopiperazino, or azepino
4 group.

1 87. The compound of claim 76, wherein R^1 is a 2,4-
2 disubstituted phenylethyl group.

1 88. The compound of claim 76, wherein R^1 is selected from
2 the group consisting of phenylethyl, 2,4-dichlorophenylethyl, 4-
3 methoxyphenylethyl, 4-phenoxyphenylethyl, 4-bromophenylethyl, 4-
4 methylphenylethyl, 4-chlorophenylethyl, 4-ethylphenylethyl, cyclohexenylethyl,
5 2-methoxyphenylethyl, 2-chlorophenylethyl, 2-fluorophenylethyl, 3-
6 methoxyphenylethyl, 3-fluorophenylethyl, thienylethyl, 4-hydroxyphenylethyl,
7 3,4-dimethoxyphenylethyl, 2-chloro-4-iodophenylethyl, 2-fluoro-4-
8 methylphenylethyl, 2-fluoro-4-chlorophenylethyl, 2-fluoro-4-bromophenylethyl,
9 2-fluoro-4-methoxyphenylethyl, 2-trifluoromethyl-4-fluorophenylethyl, 2,4-
10 difluorophenylethyl, 2,4-dimethylphenylethyl, 2,4-dimethoxyphenylethyl
11 groups, (2-pyridyl)ethyl, (3-pyridyl)ethyl, (4-pyridyl)ethyl,
12 (pyridyl)(hydroxymethyl)ethyl, (phenyl)(hydroxymethyl)ethyl, substituted and
13 unsubstituted (heteroaryl)(hydroxymethyl)ethyl, substituted and unsubstituted
14 (aryl)(hydroxymethyl)ethyl groups, substituted and unsubstituted
15 (aryl)(alkoxymethyl)ethyl, substituted and unsubstituted
16 (aryl)(aryloxymethyl)ethyl, substituted and unsubstituted
17 (aryl)(arylalkoxymethyl)ethyl, substituted and unsubstituted
18 (aryl)(heteroaryloxymethyl)ethyl, substituted and unsubstituted

19 (aryl)(heterocycloxyethyl)methyl, substituted and unsubstituted
20 (heteroaryl)(alkoxyethyl)methyl, substituted and unsubstituted
21 (heteroaryl)(aryloxyethyl)methyl, substituted and unsubstituted
22 (heteroaryl)(arylalkoxyethyl)methyl, substituted and unsubstituted
23 (heteroaryl)(heteroaryloxyethyl)methyl, and substituted and unsubstituted
24 (heteroaryl)(heterocycloxyethyl)methyl groups.

1 89. The compound of claim 76, wherein R^1 is a substituted or
2 unsubstituted alkenyl group.

1 90. A composition comprising the compound of claim 76 and
2 a pharmaceutically acceptable carrier.

1 91. A method of treating an MC4-R mediated disease,
2 comprising administering to a subject in need thereof, the compound of claim
3 76.

1 92. The method according to claim 91, wherein the disease is
2 obesity or type II diabetes.

Best Available Copy

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/16442

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 471/04, 475/04, 487/04; A61K 31/517, 31/519; A61P 3/10

US CL : 544/184, 236, 256, 257, 273, 283; 514/249, 252.02, 258, 259

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/184, 236, 256, 257, 273, 283; 514/249, 252.02, 258, 259

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE, EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2002/0193595 A1 (CHU et al.) 19 December 2002(19.12.2002), see entire document, especially formula I.	1-92

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" documents referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z"

document member of the same patent family

Date of the actual completion of the international search

04 October 2003 (04.10.2003)

Date of mailing of the international search report

09 OCT 2003

Name and mailing address of the ISA/US

Mall Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Venkataraman Balasubramanian

Telephone No. (703) 308-0196